



Cambridge IGCSE™

ADDITIONAL MATHEMATICS

0606/22

Paper 2

February/March 2023

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the February/March 2023 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **11** printed pages.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Maths-Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method marks, awarded for a valid method applied to the problem.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- B** Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation '**dep**' is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations


awrt	answers which round to
cao	correct answer only
dep	dependent
FT	follow through after error
isw	ignore subsequent working
nfwf	not from wrong working
oe	or equivalent
rot	rounded or truncated
SC	Special Case
soi	seen or implied

Question	Answer	Marks	Partial Marks
1	Correct sketch	3	B1 for correct shape, with 3 consistent maxima, 2 cusps on the x-axes and reasonable symmetry B1 for $\left(\frac{\pi}{4}, 0\right)$ and $\left(\frac{3\pi}{4}, 0\right)$ either seen on the graph or stated; must have attempted a graph of correct shape B1 for starts with (0, 4) and ends with $(\pi, 4)$ and (0, 4) either seen on the graph or stated; must have attempted a graph correct shape
2	$\frac{11x^2}{12+1-4\sqrt{3}}$	B1	
	$\frac{11x^2(13+4\sqrt{3})}{(13-4\sqrt{3})(13+4\sqrt{3})}$	M1	FT their expression of equivalent difficulty
	$\frac{11x^2(13+4\sqrt{3})}{169-48}$ or $\frac{143x^2+44\sqrt{3}x^2}{169-48}$	A1	
	$\frac{x^2(13+4\sqrt{3})}{11}$ or $\frac{13x^2+4\sqrt{3}x^2}{11}$	A1	mark final answer
	Alternative method		
	$\left(\frac{x\sqrt{11}(2\sqrt{3}+1)}{(2\sqrt{3}-1)(2\sqrt{3}+1)}\right)^2$	(M1)	
	$\left(\frac{x\sqrt{11}(2\sqrt{3}+1)}{12-1}\right)^2$ or $\left(\frac{2\sqrt{33}x+x\sqrt{11}}{12-1}\right)^2$	(A1)	
	$\frac{11x^2(12+1+4\sqrt{3})}{121}$ or $\frac{132x^2+11x^2+4\sqrt{363}x^2}{121}$	(A1)	
	$\frac{x^2(13+4\sqrt{3})}{11}$	(A1)	mark final answer

Question	Answer	Marks	Partial Marks
3	$(5x + 4)^2 * (2x - 3)^2$ soi where * is any inequality sign or =	M1	
	$21x^2 + 52x + 7 * 0$	A1	
	Critical values: $-\frac{1}{7}, -\frac{7}{3}$ soi	A1	
	$-\frac{7}{3} \leq x \leq -\frac{1}{7}$ mark final answer	A1	FT their derived critical values
	Alternative method		
	$5x + 4 * 2x - 3$ oe soi and $5x + 4 * 3 - 2x$ oe soi where * is any inequality sign or =	(M1)	
	Critical values: $-\frac{1}{7}, -\frac{7}{3}$ soi	(A2)	A1 for $-\frac{1}{7}$ or $-\frac{7}{3}$
	$-\frac{7}{3} \leq x \leq -\frac{1}{7}$ mark final answer	(A1)	FT their derived critical values
4	$[y =] \frac{1}{\operatorname{cosec} 5x} = \sin 5x$ nfwv	B1	
	$\int_0^{\frac{\pi}{5}} y \, dx = \left[-\frac{\cos 5x}{5} \right]_0^{\frac{\pi}{5}}$	B1	FT their $a \sin 5x$
	$-\frac{1}{5} \cos \left(5 \times \frac{\pi}{5} \right) - \left(-\frac{1}{5} \cos(5 \times 0) \right)$	M1	FT their $a(k \cos bx)$ where $k < 0$ or $k = \frac{1}{5}$
	$\frac{2}{5}$	A1	
5(a)	$1^3 - 2(1^2) - 19 + 20 = 0$	1	
5(b)	$(x - 1)(x^2 - x - 20)$	M2	M1 for two terms correct in the quadratic factor
	$(x - 1)(x + 4)(x - 5)$	A1	
5(c)	$e^y = 1, e^y = 5$	M1	
	$y = 0, y = \ln 5$ mark final answer	A1	1.61 or decimal equivalent for $\ln 5$ seen is A0 as calculator use not permitted

Question	Answer	Marks	Partial Marks
6(a)(i)	$\frac{1}{8}$ or 0.125	2	M1 for $64(0.5)^9$ oe
6(a)(ii)	$\frac{1023}{8}$ or 127.875	2	M1 for $\frac{64(1-0.5^{10})}{1-0.5}$ oe
6(a)(iii)	128	1	
6(b)	$\frac{20}{2}\{2a+19d\} - 400 = 2 \times \frac{10}{2}\{2a+9d\}$ oe, soi	M2	M1 for $\frac{20}{2}\{2a+19d\}$ or $\frac{10}{2}\{2a+9d\}$ soi
	$5a = a + 5d$ soi	M1	
	$d = 4$	A1	
	$a = 5$	A1	
	27 nfw	B1	must have earned all previous marks
7(a)	$\frac{d}{dx}(\cos^2 x) = -2\cos x \sin x$ soi	B1	
	Attempts the quotient rule $\frac{dy}{dx} = \frac{-2\cos x \sin x \tan x - (1 + \cos^2 x)\sec^2 x}{\tan^2 x}$	M1	FT their $\frac{d}{dx}(\cos^2 x)$
	Fully correct isw	A1	FT their $\frac{d}{dx}(\cos^2 x)$ only
	$\frac{\delta y}{h} \approx \text{their } \frac{dy}{dx} \Big _{x=\frac{\pi}{4}}$	M1	
	$\delta y \approx -4h$ cao	A1	

Question	Answer	Marks	Partial Marks
7(a)	Alternative method 1		
	$\frac{d}{dx}(\cos^3 x) = -3\cos^2 x \sin x$ soi	(B1)	
	Attempts the quotient rule: $\frac{dy}{dx} = \frac{(\sin x)(-\sin x - 3\cos^2 x \sin x) - (\cos x + \cos^3 x)\cos x}{\sin^2 x}$	(M1)	FT their $\frac{d}{dx}(\cos^3 x)$
	Fully correct isw	(A1)	FT their $\frac{d}{dx}(\cos^3 x)$ only
	$\frac{\delta y}{h} \approx \text{their } \frac{dy}{dx} \Big _{x=\frac{\pi}{4}}$	(M1)	
	$\delta y \approx -4h \text{ cao}$	(A1)	
	Alternative method 2		
	$\frac{d}{dx}\left(\frac{2}{\tan x}\right) = -2(\tan x)^{-2} \sec^2 x$	(B1)	
	Attempts the product rule $\frac{dy}{dx} = -2(\tan x)^{-2} \sec^2 x - (\sin x(-\sin x) + \cos x(\cos x))$	(M1)	FT their $\frac{d}{dx}\left(\frac{2}{\tan x}\right)$
	Fully correct isw	(A1)	FT their $\frac{d}{dx}\left(\frac{2}{\tan x}\right)$ only
	$\frac{\delta y}{h} \approx \text{their } \frac{dy}{dx} \Big _{x=\frac{\pi}{4}}$	(M1)	
	$\delta y \approx -4h \text{ cao}$	(A1)	

Question	Answer	Marks	Partial Marks
7(b)	$\frac{dy}{dx} = -3(x-3)^{-4}$ oe, soi	B1	
	$\frac{d^2y}{dx^2} = -4 \times -3(x-3)^{-5}$ oe, soi	B1	
	$\frac{(x-3)^2 + 3(x-3) - 4}{(x-3)^5}$ or $\left[\frac{x-3+3}{(x-3)^4} - \frac{4}{(x-3)^5} \right] = \frac{x(x-3) - 4}{(x-3)^5}$	M1	FT $\frac{dy}{dx} = k(x-3)^{-4}$ and $\frac{d^2y}{dx^2} = m(x-3)^{-5}$ where k and m are constants
	Correct completion to given answer: $\frac{x^2 - 3x - 4}{(x-3)^5} = \frac{(x+1)(x-4)}{(x-3)^5}$	A1	
8(a)(i)	$3 \leq x < 5$	B2	B1 for $x \geq 3$ or for $x < 5$ or for 3 and 5 in an incorrect inequality
8(a)(ii)	$x = \sqrt{5x-4}$ and rearrangement to $x^2 - 5x + 4 = 0$	B1	
	Factorises $x^2 - 5x + 4$ or solves <i>their</i> $x^2 - 5x + 4 = 0$	M1	
	$x = 4$ only, nfw	A1	
8(a)(iii)	Correct pair of graphs. 	4	B1 for correct shape for f ; may not be over correct domain but must have positive y -intercept and appear to tend to an asymptote in the 1st quadrant B1 for $(0, 3)$ and f in 1st quadrant only; must have attempted correct shape B1 for asymptote at $y = 5$; must have attempted correct shape B1 for a correct reflection of <i>their</i> f in the line $y = x$ Maximum of 3 marks if not fully correct

Question	Answer	Marks	Partial Marks
8(b)	$f^{-1}(x) = -\ln \frac{5-x}{2}$ or $f^{-1}(x) = \ln \frac{2}{5-x}$ oe	2	M1 for a complete attempt to find the inverse function with at most one sign or arithmetic error: Putting $y = f(x)$ and changing subject to x and swapping x and y or swapping x and y and changing subject to y
	Correct simplified form e.g. $\left[f^{-1}g(x) = \right] -\ln \frac{2-5x}{2(1-x)}$ or $\left[f^{-1}g(x) = \right] \ln \frac{2-2x}{2-5x}$	2	M1 FT for a correct unsimplified form of the function; FT providing of equivalent difficulty
9(a)	$6.5\left(\frac{3\pi}{8}\right) + 5.2\left(\frac{3\pi}{8}\right) + 2(6.5 - 5.2)$	M2	M1 for $6.5\left(\frac{3\pi}{8}\right)$ or $5.2\left(\frac{3\pi}{8}\right)$
	16.38 to 16.4	A1	
9(b)	[Angle $PRQ =$] 2ϕ soi	B1	
	$y = 2a \cos \phi$ oe or $y = \frac{a \sin(\pi - 2\phi)}{\sin \phi}$ oe $y^2 = a^2 + a^2 - 2a^2 \cos(\pi - 2\phi)$ oe or $y^2 = a^2 + a^2 + 2a^2 \cos(2\phi)$ oe	B1	
	Complete and correct plan soi: $\pi a^2 - \frac{1}{2}(2a \cos \phi)^2 (2\phi)$ oe or $\pi a^2 - \frac{1}{2}\left(\frac{a \sin(\pi - 2\phi)}{\sin \phi}\right)^2 (2\phi)$ oe or $\pi a^2 - \frac{1}{2}(a^2 + a^2 - 2a^2 \cos(\pi - 2\phi))(2\phi)$ oe or $\pi a^2 - \frac{1}{2}(a^2 + a^2 + 2a^2 \cos(2\phi))(2\phi)$	M1	FT their 2ϕ and their expression for y or y^2 in terms of a and ϕ
	$a^2(\pi - 4\phi \cos^2 \phi)$ or $\pi a^2 - \frac{a^2 \phi \sin^2(\pi - 2\phi)}{\sin^2 \phi}$ or $\pi a^2 - 2\phi(a^2 - a^2 \cos(\pi - 2\phi))$ or $\pi a^2 - 2\phi(a^2 + a^2 \cos 2\phi)$ oe	A1	

Question	Answer	Marks	Partial Marks
10(a)(i)	$v = 3t^2 + c$	M1	
	$v = 3t^2 - 1$	A1	
	When $t = 3$ $v = 26$	A1	
10(a)(ii)	$s = t^3 - t + c$	M1	FT $kt^2 + c$
	$s = t^3 - t - 4$	A1	
	When $t = 3$ $s = 20$	A1	
10(b)	$v = \frac{-18e^3}{e^t} + c$ oe	M1	
	$v = \frac{-18e^3}{e^t} + 44$ oe	A1	FT (<i>their</i> 26) + 18
	$s = \frac{18e^3}{e^t} + \text{their } 44t + d$ oe	M1	dep on previous M1
	$s = \frac{18e^3}{e^t} + 44t - 130$ oe, cao	A1	

Question	Answer	Marks	Partial Marks
11	Solves $\sin(4x - \pi) = 0$ oe	M1	
	$a = \frac{3\pi}{4}$	A1	
	$\frac{dy}{dx} = 4\cos(4x - \pi)$	B2	B1 for $\frac{dy}{dx} = k\cos(4x - \pi)$, where $k > 0$ or $k = -4$
	$\left[\frac{-1}{4\cos(4 \times \text{their } \frac{3\pi}{4} - \pi)} = \right] -\frac{1}{4}$	B2	FT their $a = \frac{n\pi}{4}$, n is a positive integer B1 for $\frac{-1}{\text{their } k \cos(4 \times \text{their } \frac{3\pi}{4} - \pi)}$
	$y - 0 = -\frac{1}{4}\left(x - \frac{3\pi}{4}\right)$ or $0 = -\frac{1}{4}\left(\frac{3\pi}{4}\right) + c$ oe	M1	FT their perpendicular gradient and their a
	$B\left(0, \frac{3\pi}{16}\right)$ soi	B1	
	[Exact area =] $\frac{9\pi^2}{128}$	B1	