



|  |
|--|
|  |
|--|

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

0606/13

May/June 2017

**2 hours**

Additional Materials: Electronic calculator

## READ THESE INSTRUCTIONS FIRST

DO **NOT** WRITE IN ANY BARCODES.

You are reminded of the need for clear presentation in your answers.

The total number of marks for this paper is 80.

This document consists of **15** printed pages and **1** blank page.

**Mathematical Formulae****1. ALGEBRA***Quadratic Equation*

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

*Binomial Theorem*

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

**2. TRIGONOMETRY***Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$


*Formulae for  $\triangle ABC$* 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$


$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

- 1 (a) On the Venn diagram below, shade the region which represents  $(A \cap B') \cup (C \cap B')$ . [1]



- (b) Complete the Venn diagram below to show the sets  $Y$  and  $Z$  such that  $Z \subset X \subset Y$ . [1]




- 2 Given that  $y = 3 + 4 \cos 9x$ , write down

- (i) the amplitude of  $y$ , [1]


- (ii) the period of  $y$ . [1]

- 3 (i) On the axes below, sketch the graph of  $y = 3 \sin x - 2$  for  $0^\circ \leq \theta \leq 360^\circ$ . [3]



- (ii) Given that  $0 \leq |3 \sin x - 2| \leq k$  for  $0^\circ \leq x \leq 360^\circ$ , write down the value of  $k$ . [1]

- 4 In this question, all dimensions are in centimetres.



The diagram shows an isosceles triangle  $ABC$ , where  $AB = AC$ . The point  $M$  is the mid-point of  $BC$ . Given that  $AM = 3 + 2\sqrt{5}$  and  $BC = 4 + 6\sqrt{5}$ , find, **without using a calculator**,

- (i) the area of triangle  $ABC$ , [2]

- (ii)  $\tan ABC$ , giving your answer in the form  $\frac{a + b\sqrt{5}}{c}$  where  $a$ ,  $b$  and  $c$  are positive integers. [3]

- 5 The normal to the curve  $y = \sqrt{4x + 9}$ , at the point where  $x = 4$ , meets the  $x$ - and  $y$ -axes at the points  $A$  and  $B$ . Find the coordinates of the mid-point of the line  $AB$ . [7]

6 (a) Given that  $\mathbf{A} = \begin{pmatrix} 3 & 1 \\ 2 & 4 \end{pmatrix}$ ,  $\mathbf{B} = \begin{pmatrix} 5 & 1 \\ 2 & 4 \\ -1 & 0 \end{pmatrix}$  and  $\mathbf{C} = \begin{pmatrix} -5 & 2 \\ 3 & 1 \end{pmatrix}$ , find

(i)  $\mathbf{A} + 3\mathbf{C}$ , [2]

(ii)  $\mathbf{BA}$ . [2]

(b) (i) Given that  $\mathbf{X} = \begin{pmatrix} 1 & -3 \\ 4 & -2 \end{pmatrix}$ , find  $\mathbf{X}^{-1}$ . [2]

(ii) Hence find  $\mathbf{Y}$ , such that  $\mathbf{XY} = \begin{pmatrix} 5 & -10 \\ 15 & 20 \end{pmatrix}$ . [3]

7    (a) Show that  $\frac{\tan^2 \theta + \sin^2 \theta}{\cos \theta + \sec \theta} = \tan \theta \sin \theta$ . [4]

- (b) Given that  $x = 3 \sin \phi$  and  $y = \frac{3}{\cos \phi}$ , find the numerical value of  $9y^2 - x^2y^2$ . [3]

- 8 It is given that  $p(x) = 2x^3 + ax^2 + 4x + b$ , where  $a$  and  $b$  are constants. It is given also that  $2x + 1$  is a factor of  $p(x)$  and that when  $p(x)$  is divided by  $x - 1$  there is a remainder of  $-12$ .

(i) Find the value of  $a$  and of  $b$ . [5]

(ii) Using your values of  $a$  and  $b$ , write  $p(x)$  in the form  $(2x + 1)q(x)$ , where  $q(x)$  is a quadratic expression. [2]

(iii) Hence find the exact solutions of the equation  $p(x) = 0$ . [2]

9 It is given that  $\int_{-k}^k (15e^{5x} - 5e^{-5x})dx = 6$ .

(i) Show that  $e^{5k} - e^{-5k} = 3$ . [5]

(ii) Hence, using the substitution  $y = e^{5k}$ , or otherwise, find the value of  $k$ . [3]

**10** It is given that  $y = (10x + 2)\ln(5x + 1)$ .

**(i)** Find  $\frac{dy}{dx}$ . [4]

**(ii)** Hence show that  $\int \ln(5x + 1) dx = \frac{(ax + b)}{5} \ln(5x + 1) - x + c$ , where  $a$  and  $b$  are integers and  $c$  is a constant of integration. [3]

- (iii) Hence find  $\int_0^{\frac{1}{5}} \ln(5x + 1) dx$ , giving your answer in the form  $\frac{d + \ln f}{5}$ , where  $d$  and  $f$  are integers. [2]

**11** A curve has equation  $y = 6x - x\sqrt{x}$ .

**(i)** Find the coordinates of the stationary point of the curve. [4]

**(ii)** Determine the nature of this stationary point. [2]

**(iii)** Find the approximate change in  $y$  when  $x$  increases from 4 to  $4 + h$ , where  $h$  is small. [3]

**12** A particle moves in a straight line, such that its velocity,  $v \text{ ms}^{-1}$ ,  $t$  s after passing a fixed point  $O$ , is given by  $v = 2 + 6t + 3 \sin 2t$ .

**(i)** Find the acceleration of the particle at time  $t$ . [2]

**(ii)** Hence find the smallest value of  $t$  for which the acceleration of the particle is zero. [2]

**(iii)** Find the displacement,  $x$  m from  $O$ , of the particle at time  $t$ . [5]

**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.