

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a + l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY*Identities*

$$\begin{aligned}\sin^2 A + \cos^2 A &= 1 \\ \sec^2 A &= 1 + \tan^2 A \\ \operatorname{cosec}^2 A &= 1 + \cot^2 A\end{aligned}$$


Formulae for $\triangle ABC$

$$\begin{aligned}\frac{a}{\sin A} &= \frac{b}{\sin B} = \frac{c}{\sin C} \\ a^2 &= b^2 + c^2 - 2bc \cos A \\ \Delta &= \frac{1}{2}bc \sin A\end{aligned}$$

- 1 (a) Solve the equation $\frac{|4x-5|}{7} = 1$.

[2]

(b)



The diagram shows the graph of $y = |3x + 9|$.

By drawing a suitable graph on the same diagram, solve the inequality $|3x + 9| \leq |x - 5|$. [3]

2 DO NOT USE A CALCULATOR IN THIS QUESTION.

Write the expression $\frac{\sqrt{98x^{12}}}{3+\sqrt{2}}$ in the form $(a\sqrt{b}+c)x^d$ where a , b , c and d are integers. [4]

3 (a) Differentiate $\ln(x^3 + 3x^2)$ with respect to x , simplifying your answer. [2]

(b) Hence find $\int \frac{x+2}{x(x+3)} dx$. [2]

4 The polynomial p is such that $p(x) = 2x^3 + 11x^2 + 22x + 40$.

(a) Show that $x = -4$ is a root of the equation $p(x) = 0$. [1]

(b) Factorise $p(x)$ and hence show that $p(x) = 0$ has no other real roots. [4]

- 5 (a) (i) A gardening group has 20 members. A committee of 6 members is to be selected. Anwar and Bo belong to the gardening group and at most one of them can be on the committee. How many different committees are possible? [2]

- (ii) The gate for the garden has a lock with a 6-character passcode. The passcode is to be made from


Letters	G	A	R	D	E	N					
Numbers	0	1	2	3	4	5	6	7	8	9	

No character may be used more than once in any passcode.

Find the number of possible passcodes that have 4 letters followed by 2 numbers. [2]

(b) (i) Given that $n \geq 4$, show that $(n-3) \times {}^nC_3 = 4 \times {}^nC_4$. [2]

(ii) Given that ${}^nC_3 = 5n$, where $n \geq 3$, show that n satisfies the equation $n^2 - 3n - 28 = 0$.
Hence find the value of n . [4]



The diagram shows the curve $y = 5e^{2x} - 3$. The curve meets the y -axis at the point A . The tangent to the curve at A meets the x -axis at the point B . Find the length of AB . [6]

- 7 Variables x and y are such that $y = \frac{4x^3 + 2 \sin 8x}{1-x}$. Use differentiation to find the approximate change in y as x increases from 0.1 to $0.1 + h$, where h is small. [6]

- 8 (a) The functions f and g are defined by

$$f(x) = \sec x \quad \text{for } \frac{\pi}{2} < x < \frac{3\pi}{2}$$

$$g(x) = 3(x^2 - 1) \quad \text{for all real } x.$$


- (i) Find the range of f . [1]

- (ii) Solve the equation $f^{-1}(x) = \frac{2\pi}{3}$. [3]

- (iii) Given that gf exists, state the domain of gf . [1]

- (iv) Solve the equation $gf(x) = 1$. [5]


- (b) The function h is defined by $h(x) = \ln(4-x)$ for $x < 4$. Sketch the graph of $y = h(x)$ and hence sketch the graph of $y = h^{-1}(x)$. Show the position of any asymptotes and any points of intersection with the coordinate axes. [4]



9 (a) Show that $\int_1^8 \frac{x+4}{\sqrt[3]{x}} dx = 36.6$.

[3]

(b)



The diagram shows part of the line $10y = 7 - 3x$ and part of the curve $y = \frac{1}{3x+4}$.

The line and curve intersect at the point A. Verify that the y-coordinate of A is 0.1 and calculate the area of the shaded region.

[8]

Continuation of working space for Question 9(b).

- 10** An arithmetic progression, A , has first term a and common difference d .
The 2nd, 14th and 17th terms of A form the first three terms of a convergent geometric progression, G , with common ratio r .

(a) (i) Given that $d \neq 0$, find two expressions for r in terms of a and d and hence show that $a = -17d$.
[6]

(ii) Find the value of r . [2]

- (b) The first term of the geometric progression, G , is q and the sum to infinity is $\frac{256}{3}$.

Find the sum of the first 20 terms of the **arithmetic** progression, A .

[7]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.