



Cambridge International Examinations
Cambridge International General Certificate of Secondary Education

BIOLOGY

0610/53

Paper 5 Practical Test

May/June 2017

MARK SCHEME

Maximum Mark: 40

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

© IGCSE is a registered trademark.

This syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **6** printed pages.

Mark schemes will use these abbreviations

- ; separates marking points
- / alternatives
- **I** **I**
- **R** reject
- **A** **A** (for answers correctly cued by the question, or guidance for examiners)
- AW alternative wording (where responses vary more than usual)
- AVP any valid point
- **ecf** credit a correct statement / calculation that follows a previous wrong response
- **ora** or reverse argument
- () the word / phrase in brackets is not required, but sets the context
- underline actual word given must be used by candidate (grammatical variants excepted)
- max indicates the maximum number of marks that can be given

Question	Answer	Marks	Guidance
1(a)	table with 2 columns ; column 1 heading 'vegetable extract' column 2 headed volume of iodine and column 2 units – cm ³ ; correct trend ;	3	
1(b)	to allow iodine to change colour ;	1	A as an indicator
1(c)	volume of vegetable extract ; volume / concentration of starch solution ; concentration of iodine ; temperature; mixing time;	2	I amount R volume iodine solution

Question	Answer	Marks	Guidance
1(d)	source of error ;;	4	improvement must relate to given error A subjective colour change
	contamination		
	overshoot of end-point / adding too much iodine / many drops		
	determination of end-point		
	change in vitamin C with time		
	no repeats		
	AVP e.g. difficult reading scale coloured vegetable extracts		
	improvement ;;		
	washing all apparatus / use new syringes		
	add smaller quantities of iodine		
	allow sufficient time for colour to change / use colorimeter / colour standard		
	test same time after extraction for each		
	repeat each concentration		
	AVP e.g. use burette		
1(e)(i)	L: 25.00 ; N: 62.5 ; correct number of decimal places on both ;	3	
1(e)(ii)	axes labelled and units; even scale to fill more than half of printed grid ; plot three / four points correctly ; line of best fit / trend line ;	4	ecf candidate result for 1(e)(i)

Question	Answer	Marks	Guidance
1(e)(iii)	mark volume of iodine used on (y axis of) graph / extend horizontally and extend line vertically from plotted point to x axis ; correct reading from graph on answer line ;	2	
1(f)	range of temperatures ; values for temperatures stated ; time at each temperature ; use of water bath / named method ; description of extracting juice ; detail of use of iodine drops / volume / addition of starch for end point ;; at least two repeats ; (controlled variables) heating time / same type of vegetable / all samples from same vegetable ; relevant reference to safety ;	6	minimum of three at least one above 50

Question	Answer	Marks	Guidance
2(a)	<i>any four from:</i> drawing with clear outline ; scaled to fit more than half the space ; shape 5 / 6 sides for both ; detail showing 3 / 4 layers with no shading and no cells ;	4	
2(b)(i)	length of PQ = 80 mm ; (x)64 ;	3	$\pm 1\text{mm}$ $(80) \div 1.25$
2(b)(ii)	plane of section ; AW magnification ; number of villi different;	2	
2(c)(i)	A: 3 B: 9 C: 11 ;	1	A 2 instead of 3 for A 3 correct answers = 1 mark
2(c)(ii)	30 °C ; has highest rate of reaction / AW ;	2	
2(c)(iii)	it is much higher / different than trial 1 and 3 / AW ;	1	
2(c)(iv)	(IV) temperature ; (DV) rate of reaction ;	2	