



Cambridge O Level

ADDITIONAL MATHEMATICS

4037/21

Paper 2

May/June 2023

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2023 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Maths-Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark


- M** Method marks, awarded for a valid method applied to the problem.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- B** Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation ‘**dep**’ is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations

- awrt answers which round to
- cao correct answer only
- dep dependent
- FT follow through after error
- isw ignore subsequent working
- nfwf not from wrong working
- oe or equivalent
- rot rounded or truncated
- SC Special Case
- soi seen or implied


Question	Answer	Marks	Partial Marks
1	$\lg y = 2\sqrt{x} + 3$ OR $\lg b = 2$ and $\lg A = 3$	B2	B1 for $\lg y = \left(\frac{8-5}{2.5-1}\right)\sqrt{x} + c$ soi or $\lg y = m\sqrt{x} + 3$ soi OR $\lg b = \frac{8-5}{2.5-1}$ or $\lg A = 3$ soi
	$y = 10^{2\sqrt{x}+3}$ or $\lg \frac{y}{10^3} = 2\sqrt{x}$ OR $b = 100$ and $A = 1000$	M1	FT their m and c
	$y = 10^3 \times 100^{\sqrt{x}}$ oe mark final answer	A1	

Question	Answer	Marks	Partial Marks
2(a)	Correct curve 	3	B2 for correct cosine shape over 2 cycles with midline at $y = 2$ and consistent amplitude or B1 for attempt at cosine shape over 2 cycles with consistent amplitude B1 for a consistent amplitude of 2; must have attempted correct shape Maximum of 2 marks if not fully correct
2(b)	4	1	
2(c)	60°	1	
3(a)	$a = 2, b = 3, c = -2$	2	B1 for any two correct
3(b)	$-3 \leq x \leq -0.5$ or $x \geq 1$	3	B1 for the critical values $-3, -0.5, 1$ B1 for $-3 \leq x \leq -0.5$ B1 for $x \geq 1$
4(a)	$(2y - 1) \log 5 = \log 6 + y \log 3$ oe or $2y \log 5 = \log 30 + y \log 3$ oe OR [rearranges $\frac{5^{2y}}{5} = 6 \times 3^y$ and collects powers to a single power in y] $\left(\frac{5^2}{3}\right)^y = 30$ oe	M1	
	Collects terms and factorises: $y(2 \log 5 - \log 3) = \log 6 + \log 5$ oe or $y(2 \log 5 - \log 3) = \log 30$ oe OR takes logs $y = \log_{\frac{25}{3}} 30$ oe or $y \log \left(\frac{5^2}{3}\right) = \log 30$ oe	M1	FT if of equivalent difficulty
	1.604	A1	cao

Question	Answer	Marks	Partial Marks
4(b)	$e^{4x} - 4e^{2x} + 3 = 0$ or $(e^{2x})^2 - 4e^{2x} + 3 = 0$ oe	M1	condone one error
	Factorises: $(e^{2x} - 1)(e^{2x} - 3)$ oe or solves $e^{4x} - 4e^{2x} + 3 = 0$ oe	M1	FT $a(e^{2x})^2 + be^{2x} + c [= 0]$
	$e^{2x} = 1, e^{2x} = 3$	A1	
	$x = 0, x = \frac{1}{2} \ln 3$ or exact equivalent	A1	
5	$\frac{dV}{dr} = 4\pi r^2$ oe and $\left. \frac{dV}{dr} \right _{r=6} = 144\pi$	B1	
	$\frac{dr}{dt} = \frac{dV}{dt} \times \frac{dr}{dV}$ soi	B1	Not if chain rule for $\frac{dt}{dr}$ unless answer is inverted
	$\frac{24}{\text{their } 144\pi}$	M1	<i>their</i> 144π must come from an attempt at differentiation
	0.0531 or 0.05305[16....] rot to 4 or more sig figs	A1	
6(a)	$3\begin{pmatrix} x-8 \\ y-5 \end{pmatrix} = 2\begin{pmatrix} x-4 \\ y-7 \end{pmatrix}$ oe OR $\overrightarrow{QR} = \begin{pmatrix} x-8 \\ y-5 \end{pmatrix}$ and $\overrightarrow{PR} = \begin{pmatrix} x-4 \\ y-7 \end{pmatrix}$ and $3x - 24 = 2x - 8$ and $3y - 15 = 2y - 14$	M1	
	$x = 16$	A1	dep on vector method
	$y = 1$	A1	dep on vector method

Question	Answer	Marks	Partial Marks
6(b)(i)	$\mathbf{a} = -2.5\mathbf{i} - \frac{5\sqrt{3}}{2}\mathbf{j}$ isw	B1	
	$\mathbf{c} = -5\mathbf{i} + 5\sqrt{3}\mathbf{j}$ isw	B1	
6(b)(ii)	$\mathbf{b} = (-5 + 2.5)\mathbf{i} + (5\sqrt{3} + 2.5\sqrt{3})\mathbf{j}$ oe soi	B1	
	$r = \sqrt{(-2.5)^2 + (7.5\sqrt{3})^2}$	M1	FT <i>their b</i> of the form $x\mathbf{i} + y\mathbf{j}$ providing neither component is zero
	$[r =] 13.2$ or 13.22875... rot to 4 or more sf	A1	dep on B1
	$\tan \alpha = \left(\frac{7.5\sqrt{3}}{2.5} \right)$ oe or awrt 79.1 or $\tan \beta = \left(\frac{2.5}{7.5\sqrt{3}} \right)$ oe or awrt 10.9	M1	FT <i>their b</i> of the form $x\mathbf{i} + y\mathbf{j}$
	349[.106...] rot to 3 or more sf	A1	dep on B1
	Alternative method		
	$[r^2 =] 10^2 + 5^2 - 2 \times 10 \times 5 \times \cos 120^\circ$	(M1)	
	$[r =] 13.2$ or 13.22875... rot to 4 or more sf	(A1)	
	$\frac{\sin \theta}{10} = \frac{\sin \text{their } 120}{\text{their } 5\sqrt{7}}$ or $\frac{\sin \phi}{5} = \frac{\sin \text{their } 120}{\text{their } 5\sqrt{7}}$	(M1)	FT consistent use of <i>their</i> 120 and <i>their</i> r
	$[\theta =]$ awrt 40.9 or $[\phi =]$ awrt 19.1	(A1)	
	349[.106...] rot to 3 or more sf	(A1)	

Question	Answer	Marks	Partial Marks
7(a)	$\left[\frac{6x^2}{2} - \frac{x^3}{3} \right]_0^5$	B1	
	Area under line: $0.5 \times 5 \times 5$ oe	B1	
	Fully actioned correct plan: $3(25) - \frac{5^3}{3} - \left(3(0) - \frac{0^3}{3} \right) - 0.5 \times 5 \times 5$ oe	M1	
	$\frac{125}{6}$ oe isw	A1	dep on all previous marks
	Alternative method		
	$\int_0^5 (5x - x^2) dx$	(B1)	
	$\left[\frac{5x^2}{2} - \frac{x^3}{3} \right]_0^5$	(B1)	
	Correct use of correct limits $2.5(25) - \frac{5^3}{3} - \left(2.5(0) - \frac{0^3}{3} \right)$	(M1)	
	$\frac{125}{6}$ oe isw	(A1)	dep on all previous marks
7(b)(i)	$\frac{(2x-6)^{-2}}{-2 \times 2} + \sin x (+c)$ oe, isw	3	B1 for $\sin x$ B2 for $\frac{(2x-6)^{-2}}{-2 \times 2}$ or B1 for $\frac{(2x-6)^{-2}}{-2}$ soi
7(b)(ii)	$\frac{x^7}{2} + x^3 + \frac{1}{2x}$ oe	B1	
	$\frac{x^8}{16} + \frac{x^4}{4} + \frac{1}{2} \ln x (+c)$ oe or $\frac{x^8}{16} + \frac{x^4}{4} + \frac{1}{2} \ln 2x (+c)$ oe	B2	B1 for any two correct

Question	Answer	Marks	Partial Marks
8(a)(i)	[Domain f^{-1}] $0 \leq x \leq 2.25$ oe	B2	B1 for either end correct or for 0 and 2.25 in an incorrect inequality
	[Range f^{-1}] $0 \leq f^{-1} \leq 3$	B1	
8(a)(ii)	$x = 1.6$ oe or $x = 0$	2	B1 for each
8(a)(iii)		2	B1 for attempt at correct graph of inverse function drawn over correct domain soi B1 for correct shape with intersection in approximately correct location
8(b)(i)	For a complete method to find the inverse, including changing the subject and swapping the variables	M1	
	$[g^{-1}(x) =] \sqrt[3]{\frac{x^3 - 3}{8}}$ oe mark final answer	A1	
8(b)(ii)	$[k =] 0$	1	
8(b)(iii)	$\sqrt[3]{8e^{12x} + 3}$ mark final answer	1	

Question	Answer	Marks	Partial Marks
9(a)	$\frac{1}{2} \times 24^2 \times \theta = 432$	M1	
	$\theta = \frac{3}{2}$ rads soi	A1	
	$24 \times \textit{their } \theta$	M1	
	36 cao	A1	
	Alternative method		
	$s = r\theta$ soi and $\frac{1}{2} \times r \times s = 432$	(B1)	
	$\frac{1}{2} \times 24 \times s = 432$	(M1)	
	$s = \frac{432 \times 2}{24}$ oe	(M1)	
	$[s =] 36$	(A1)	
9(b)(i)	$[OB =] 2y \cos \alpha$ oe	B1	
9(b)(ii)	$\frac{(\textit{their } 2y \cos \alpha) \times y \sin \alpha}{2}$ $-\frac{1}{2} \times (\textit{their } y \cos \alpha)^2 \times \alpha$ oe	M2	M1 for either area
	correct completion to $\frac{y^2}{2} \cos \alpha (2 \sin \alpha - \alpha \cos \alpha)$	A1	

Question	Answer	Marks	Partial Marks
10	[Term independent of x :] ${}^9C_3 \times a^6 \times b^3$ or $84 \times a^6 \times b^3$	B1	
	$a^6 b^3 = \frac{-145152}{84}$	M1	dep on B1
	$(a^2 b)^3 = -1728$ leading to $a^2 b = -12$ or $a^2 b = \sqrt[3]{-1728} = -12$	A1	
	${}^9C_1 \times a^8 \times b$ or $9 \times a^8 \times b$	B1	
	Correctly solves correct equations simultaneously $9 \times a^8 \times b = -6912$ and $a^2 b = -12$ as far as $a^6 = \dots$ or $b^3 = \dots$	M1	Must be solving correct equations
	$a = 2, b = -3$ and no other values nfw	B2	B1 for each nfw dep on previous B1B1
11	Eliminates one variable $(k - 3y)^2 + y^2 + 2y - 9 = 0$	M1	
	$10y^2 + (2 - 6k)y + (k^2 - 9) = 0$ soi	A1	
	Uses $b^2 - 4ac \neq 0$ with <i>their</i> 3-term quadratic: $(2 - 6k)^2 - 4(10)(k^2 - 9)$ [*0]	M1	* can be = or any inequality sign
	$-4k^2 - 24k + 364$ [*0]	A1	
	Factorises $-4k^2 - 24k + 364$ or solves <i>their</i> $-4k^2 - 24k + 364 = 0$	M1	
	$k = 7$ only	A1	
	Uses <i>their</i> k in $10y^2 + (2 - 6k)y + (k^2 - 9) = 0$ oe	M1	
	$y = 2$ only	A1	
	$x = 1$ only	A1	