



8 5 5 5 5 2 5 1 7

--

--	--	--	--	--

--	--	--	--

9231/31

May/June 2021

1 hour 30 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 ms^{-2} .

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

BLANK PAGE

- 1 A particle P of mass 1 kg is moving along a straight line against a resistive force of magnitude $\frac{10\sqrt{v}}{(t+1)^2}\text{ N}$, where $v\text{ ms}^{-1}$ is the speed of P at time $t\text{ s}$. When $t = 0$, $v = 25$.

Find an expression for v in terms of t .

[5]

[illegible]

- 2** A hollow hemispherical bowl of radius a has a smooth inner surface and is fixed with its axis vertical. A particle P of mass m moves in horizontal circles on the inner surface of the bowl, at a height x above the lowest point of the bowl. The speed of P is $\sqrt{\frac{8}{3}ga}$.

Find x in terms of a .

[6]

[illegible]

- 3** One end of a light elastic string, of natural length a and modulus of elasticity kmg , is attached to a fixed point A . The other end of the string is attached to a particle P of mass $4m$. The particle P hangs in equilibrium a distance x vertically below A .

(a) Show that $k = \frac{4a}{x-a}$. [1]

.....

.....


.....

.....

An additional particle, of mass $2m$, is now attached to P and the combined particle is released from rest at the original equilibrium position of P . When the combined particle has descended a distance $\frac{1}{3}a$, its speed is $\frac{1}{3}\sqrt{ga}$.

(b) Find x in terms of a . [6]

[illegible]



(a) Show that the distance of the centre of mass of the combined solid from the base of the cylinder is $\frac{h(k^2 + 4k + 6)}{4(3 + k)}$. [4]

[illegible]

The solid is placed on a plane that is inclined to the horizontal at an angle θ . The base of the cylinder is in contact with the plane. The plane is sufficiently rough to prevent sliding. It is given that $3h = 2r$ and that the solid is on the point of toppling when $\tan \theta = \frac{4}{3}$.

(b) Find the value of k .

[3]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.


- 5 A particle P of mass m is attached to one end of a light inextensible string of length a . The other end of the string is attached to a fixed point O . The particle completes vertical circles with centre O . The points A and B are on the path of P , both on the same side of the vertical through O . OA makes an angle θ with the downward vertical through O and OB makes an angle θ with the upward vertical through O .

The speed of P when it is at A is u and the speed of P when it is at B is $\sqrt{ag}$. The tensions in the string at A and B are T_A and T_B respectively. It is given that $T_A = 7T_B$.

Find the value of θ and find an expression for u in terms of a and g .

[8]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.



-
- This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

The total kinetic energy of the spheres after the collision is $\frac{3}{4}mu^2$.

(b) Find the value of α .

[4]

[illegible]

- 7 A particle P is projected from a point O on a horizontal plane and moves freely under gravity. The initial velocity of P is 100 ms^{-1} at an angle θ above the horizontal, where $\tan \theta = \frac{4}{3}$. The two times at which P 's height above the plane is $H \text{ m}$ differ by 10 s .

(a) Find the value of H .

[5]

[illegible]

(b) Find the magnitude and direction of the velocity of P one second before it strikes the plane. [4]

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

[illegible]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.