



CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

9231/11

October/November 2023

2 hours

You will need: List of formulae (MF19)

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

- 1 (a) By considering $(r+1)^2 - r^2$, use the method of differences to prove that

$$\sum_{r=1}^n r = \frac{1}{2}n(n+1). \quad [4]$$

This image shows a full page of a handwriting practice worksheet. It consists of multiple rows of horizontal dotted lines spaced evenly down the page, providing a guide for letter height and placement. The background is plain white, and there are no other markings or text present.

[3]

[illegible]

2 Prove by mathematical induction that, for all positive integers n ,

$$1 + 2x + 3x^2 + \dots + nx^{n-1} = \frac{1 - (n+1)x^n + nx^{n+1}}{(1-x)^2}. \quad [6]$$

[illegible]

- 3** The quartic equation $x^4 + bx^3 + cx^2 + dx - 2 = 0$ has roots $\alpha, \beta, \gamma, \delta$. It is given that

$$\alpha + \beta + \gamma + \delta = 3, \quad \alpha^2 + \beta^2 + \gamma^2 + \delta^2 = 5, \quad \alpha^{-1} + \beta^{-1} + \gamma^{-1} + \delta^{-1} = 6.$$

- (a) Find the values of b , c and d . [6]

[illegible]

- (b) Given also that $\alpha^3 + \beta^3 + \gamma^3 + \delta^3 = -27$, find the value of $\alpha^4 + \beta^4 + \gamma^4 + \delta^4$. [2]

[illegible]

4 The lines l_1 and l_2 have equations

$$\mathbf{r} = -2\mathbf{i} - 3\mathbf{j} - 5\mathbf{k} + \lambda(-4\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}) \quad \text{and} \quad \mathbf{r} = 2\mathbf{i} - 2\mathbf{j} + 3\mathbf{k} + \mu(2\mathbf{i} - 3\mathbf{j} + \mathbf{k})$$

respectively.

(a) Find the shortest distance between l_1 and l_2 . [5]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

The plane Π contains l_1 and the point with position vector $-\mathbf{i} - 3\mathbf{j} - 4\mathbf{k}$.

- (b)** Find an equation of Π , giving your answer in the form $ax + by + cz = d$. [4]

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dotted lines, providing a guide for letter height and placement. The lines are evenly spaced across the entire page, leaving ample room for writing practice. There is no text or other markings on the page.

- 5** Let k be a constant. The matrices $\mathbf{A}$, $\mathbf{B}$ and $\mathbf{C}$ are given by

$$\mathbf{A} = \begin{pmatrix} 1 & k & 3 \\ 2 & 1 & 3 \\ 3 & 2 & 5 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 0 & -2 \\ -1 & 3 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \mathbf{C} = \begin{pmatrix} -2 & -1 & 1 \\ 1 & 1 & 3 \end{pmatrix}.$$

It is given that $\mathbf{A}$ is singular.

- (a) Show that $\mathbf{CAB} = \begin{pmatrix} 3 & -7 \\ -9 & 3 \end{pmatrix}$. [5]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- (b) Find the equations of the invariant lines, through the origin, of the transformation in the $x-y$ plane represented by **CAB**. [5]

[illegible]

- **D** represents an enlargement, centre the origin.
- **E** represents a stretch parallel to the x -axis.
- **F** represents a reflection in the line $y = x$.

[5]

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- 6 (a) Show that the curve with Cartesian equation

$$\left(x - \frac{1}{2}\right)^2 + y^2 = \frac{1}{4}$$

has polar equation $r = \cos \theta$. [3]

.....

.....

.....

.....

.....

The curves C_1 and C_2 have polar equations

$$r = \cos \theta \quad \text{and} \quad r = \sin 2\theta$$

respectively, where $0 \leq \theta \leq \frac{1}{2}\pi$. The curves C_1 and C_2 intersect at the pole and at another point P .

- (b) Find the polar coordinates of P . [3]

.....

.....

.....

.....

- (c) In a single diagram sketch C_1 and C_2 , clearly identifying each curve, and mark the point P . [3]

- (d)** The region R is enclosed by C_1 and C_2 and includes the line OP .

Find, in exact form, the area of R .

[6]

This image shows a full page of a handwriting practice worksheet. It consists of multiple rows of horizontal dotted lines spaced evenly down the page, providing a guide for letter height and placement. The background is plain white, and there are no other markings or text present.

- 7 The curve C has equation $y = f(x)$, where $f(x) = \frac{x^2 + 2}{x^2 - x - 2}$.

(a) Find the equations of the asymptotes of C .

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the coordinates of any stationary points on C , giving your answers correct to 1 decimal place.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Sketch C , stating the coordinates of any intersections with the axes. [3]

-
- (d) Sketch the curve with equation $y = \frac{1}{f(x)}$. [2]

- (e) Find the set of values for which $\frac{1}{f(x)} < f(x)$. [4]

[illegible]

[illegible]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.