



Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

FURTHER MATHEMATICS

9231/21

Paper 2 Further Pure Mathematics 2

October/November 2025

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

BLANK PAGE





1 Find the values of k for which the system of equations

$$x - y + z = 0,$$

$$x + ky + 3z = 0,$$

$$x + 2y + kz = 0,$$

does not have a unique solution.

[3]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.



2 The curve C has parametric equations

$$x = \sinh t \quad \text{and} \quad y = t + \cosh t.$$

(a) Find $\frac{dy}{dx}$ in terms of t . [2]

This image shows a single sheet of white paper with ten horizontal dashed lines, typical of primary-ruled notebook paper. The lines are evenly spaced and extend across the width of the page. There is no handwriting or other markings on the paper.

(b) Show that $\frac{d^2y}{dx^2} = \frac{1 - \sinh t}{\cosh^3 t}$. [4]

[illegible]



DO NOT WRITE IN THIS MARGIN

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the Maclaurin’s series for y in terms of x up to and including the term in x^2 . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





3 The integral I_n is defined by $I_n = \int_0^1 (1+x^2)^n dx$.

(a) By considering $\frac{d}{dx}(x(1+x^2)^n)$, or otherwise, show that

$$(2n+1)I_n = 2^n + 2nI_{n-1}. \quad [5]$$

[illegible]



(b) Find the exact value of I_{-2} .

[4]

This image shows a full page of primary-ruled paper. It features approximately 20 horizontal dotted lines spaced evenly down the page, providing a guide for handwriting practice. The paper is otherwise blank, with no margins, text, or other markings.




4 Find the particular solution of the differential equation

$$5 \frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + y = x^2 + 5x + 3,$$

given that, when $x = 0$, $y = \frac{dy}{dx} = 0$. [10]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting or typing. There are no margins, text, or other markings on the page.

[illegible]



The diagram shows the curve with equation $y = \frac{1}{3}x^3 + x$ for $0 \leq x \leq 1$, together with a set of n rectangles of width $\frac{1}{n}$.

- (a) By considering the sum of the areas of these rectangles, show that $\int_0^1 \left(\frac{1}{3}x^3 + x \right) dx < U_n$, where

$$U_n = \frac{1}{12} \left(1 + \frac{1}{n} \right) \left(7 + \frac{1}{n} \right). \quad [5]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





- (b) Use a similar method to find, in terms of n , a lower bound L_n for $\int_0^1 \left(\frac{1}{3}x^3 + x \right) dx$. [4]

- (c) Show that $\lim_{n \rightarrow \infty} (U_n - L_n) = 0$. [2]





6 The matrix $\mathbf{P}$ has non-zero eigenvalues and is given by

$$\mathbf{P} = \begin{pmatrix} a & 1 & 1 \\ 0 & 2a & -1 \\ 0 & 0 & -3a \end{pmatrix}.$$

(a) State, in terms of a , the eigenvalues of $\mathbf{P}$. [1]

.....

(b) (i) Find $\mathbf{P}^2$ in terms of a . [1]

[illegible]

(ii) Use the characteristic equation of $\mathbf{P}$ to find $\mathbf{P}^{-1}$ in terms of a . [3]

[illegible]



The 3×3 matrix $\mathbf{A}$ has eigenvalues 1, 2, 3 with corresponding eigenvectors

$$\begin{pmatrix} a \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 2a \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ -3a \end{pmatrix},$$

respectively.

(c) Find $\mathbf{A}$ in terms of a .

[5]

[illegible]



- 7 (a) Show that $\frac{d}{dx}(\tanh^{-1}x) = \frac{1}{1-x^2}$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find the solution of the differential equation

$$x \frac{dy}{dx} - y = x^2 \tanh^{-1}x,$$

for $0 < x < 1$, given that $y = 0$ when $x = \frac{1}{2}$. Give your answer in an exact form.

[9]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





This image shows a full page of a handwriting practice worksheet. It consists of approximately 20 horizontal rows. Each row is defined by two parallel dotted lines, creating a series of uniform gaps for letter height. The entire page is otherwise blank, with no margins, text, or other markings.





- 8** (a) State the sum of the series $z + z^2 + \dots + z^n$, for $z \neq 1$. [1]

.....

.....

- (b)** By letting $z = \frac{1}{2}(\cos \theta + i \sin \theta)$ use de Moivre's theorem to deduce that

$$\sum_{m=1}^n \left(\frac{1}{2}\right)^m \sin m\theta = \frac{\left(\frac{1}{2}\right)^{n+2} \sin n\theta - \left(\frac{1}{2}\right)^{n+1} \sin(n+1)\theta + \frac{1}{2} \sin \theta}{\frac{5}{4} - \cos \theta}. \quad [6]$$

[illegible]



- (c) Use the result in (b) to find $\sum_{m=1}^n \left(\frac{1}{2}\right)^m m \cos m\theta$ in terms of n and θ . [You do not need to simplify your answer.] [3]

- (d) Hence find $\sum_{m=1}^{\infty} \left(\frac{1}{2}\right)^m m \cos m\theta$ in terms of $\cos \theta$. [You may assume that $\frac{n}{2^n} \rightarrow 0$ as $n \rightarrow \infty$.] [2]





Additional page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

Lined area for writing answers, consisting of multiple horizontal dotted lines.







Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

