



CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

9701/22

February/March 2023

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **16** pages. Any blank pages are indicated.

- 1 The Pauling electronegativity values of elements can be used to predict the chemical properties of compounds.

Use the information in Table 1.1 to answer the following questions.

Table 1.1

element	H	Li	C	O	S
Pauling electronegativity value	2.1	1.0	2.5	3.5	2.6
first ionisation energy/ kJ mol^{-1}	1310	519	1090	1310	1000
second ionisation energy/ kJ mol^{-1}	—	7300	2350	3390	2260

- (a) (i) Define electronegativity.

.....
 [1]

- (ii) O and S are in Group 16.

Explain the difference in the Pauling electronegativity values of O and S.

.....

 [2]

- (b) (i) LiH is an ionic compound.

Draw a dot-and-cross diagram of LiH.

Include **all** electrons.

[2]

- (ii) Suggest the shape of a molecule of H_2S .

..... [1]

(c) (i) Write an equation that represents the first ionisation energy of H.

..... [1]

(ii) Explain why there is no information given in Table 1.1 for the second ionisation energy of H.

..... [1]

(iii) Give the full electronic configuration of $S^{2+}(g)$.

..... [1]

(d) CO_2 and SO_2 are acidic gases.

(i) Write an equation for the reaction of SO_2 with H_2O .

..... [1]

(ii) Write an equation for the reaction of SO_2 with $NaOH$.

..... [1]

(iii) Construct an equation for the reaction of CO_2 with $Mg(OH)_2$.

..... [1]

- (e) (i) Complete Table 1.2 by placing a tick (✓) to show which of the compounds have molecules with an overall dipole moment.

Table 1.2

compound	O=C=O	O=S=O	S=C=S	S=C=O
overall dipole moment				

[2]

- (ii) At 150 °C and 103 kPa, all of the compounds listed in Table 1.2 are gases.

Under these conditions, 0.284 g of one of the compounds occupies a volume of 127 cm³.

Use this information to calculate the M_r of the compound. Hence, identify the compound from those given in Table 1.2.

Show your working.

M_r = identity of compound = [3]

[Total: 17]

BLANK PAGE

2 The Group 2 elements Mg to Ba are all silvery-white reactive metals.

- (a) (i) Draw a labelled diagram to show the bonding and structure of the Group 2 metals at room temperature.

[2]

- (ii) Explain why Mg has a higher electrical conductivity than Na.

.....
..... [1]

- (b) Write an equation for the reaction of magnesium with cold water.

..... [1]

- (c) Identify a single reagent that can be used to distinguish separate samples of dilute $\text{Mg}(\text{NO}_3)_2(\text{aq})$ and dilute $\text{Ba}(\text{NO}_3)_2(\text{aq})$.

Explain your answer.

reagent

explanation

.....
[2]

- (d) (i) Describe what is observed when $\text{SrI}_2(\text{aq})$ reacts with concentrated sulfuric acid.

.....
..... [2]

- (ii) Compound **X**, an anhydrous Group 2 bromide, is dissolved in water and titrated against aqueous silver nitrate.

A solution containing 0.250 g of **X** requires 33.65 cm^3 of $0.0500 \text{ mol dm}^{-3} \text{ AgNO}_3(\text{aq})$ for complete reaction.

Identify **X**.

Show your working.

X = [3]

[Total: 11]

- 3 Alkenes undergo an addition reaction with a 1:1 mixture of CO and H₂ to form aldehydes.

Fig. 3.1 shows the reaction of propene with a 1:1 mixture of CO and H₂.




Fig. 3.1

- (a) (i) Define addition reaction.

.....
 [1]

- (ii) Aldehydes **A** and **B** are structural isomers.

State the type of structural isomerism shown by **A** and **B**.

..... [1]

- (iii) Name **A**.

..... [1]

- (iv) The complete reaction of propene with a 1:1 mixture of CO and H₂ produces **A** and **B** only. The product mixture contains 96% **A** and 4% **B**.

Calculate the mass of **A** produced in this reaction when 5.00×10^3 kg of propene is used.

mass of **A** = kg [1]

(b) **A** and **B** show reactions typical of aliphatic aldehydes.

- (i) **A** undergoes a nucleophilic addition reaction with a mixture of HCN and KCN, forming compound **C**.

Complete the diagram to show the mechanism for this reaction.

Include charges, dipoles, lone pairs of electrons and curly arrows, as appropriate.

Draw the structure of the organic intermediate.



[4]

- (ii) Table 3.1 shows information about three experiments involving **B**.

Complete Table 3.1.

Table 3.1

experiment	reagents	observation with B
1		solution turns from orange to green
2		a silver mirror forms on the sides of the reaction vessel
3	$\text{Br}_2(\text{aq})$	

[3]

- (iii) **B**, $\text{C}_4\text{H}_8\text{O}$, is oxidised by acidified potassium manganate(VII).

Complete the equation for this reaction. Use [O] to represent one atom of oxygen from the oxidising agent.



- (iv) **C** is a chiral molecule.

Circle any chiral centres in the structure of **C** shown in Fig. 3.2.




Fig. 3.2

[1]

- (c) When propene reacts with CO and an excess of H_2 , an alkane and a mixture of alcohols are formed instead. The alcohols are isomers of each other.


Suggest the molecular formulae of the alkane and the alcohols that are formed under these conditions.

molecular formula of alkane

molecular formula of alcohols

[2]

- (d) The reaction of ethene, C_2H_4 , with a 1:1 mixture of CO and H_2 is shown in equation 1.



At atmospheric pressure a cobalt-based catalyst is used in this reaction.

- (i) State and explain the effect of using a catalyst on this reaction.

.....

.....

.....

..... [2]

- (ii) Explain why the yield of $\text{CH}_3\text{CH}_2\text{CHO}(\text{g})$ increases when the overall pressure of the reaction mixture is increased.

.....
 [1]

- (iii) Use the information in Table 3.2 to calculate the enthalpy change, ΔH_r , of the reaction in equation 1.




Table 3.2

compound	enthalpy change of formation, $\Delta H_f / \text{kJ mol}^{-1}$
$\text{C}_2\text{H}_4(\text{g})$	+52
$\text{CO}(\text{g})$	−111
$\text{CH}_3\text{CH}_2\text{CHO}(\text{g})$	−187

$$\Delta H_r = \dots\dots\dots \text{kJ mol}^{-1} \quad [2]$$

- (iv) The reaction mixture is cooled to collect $\text{CH}_3\text{CH}_2\text{CHO}$ as a liquid.

Identify all types of van der Waals' forces that are present between molecules of $\text{CH}_3\text{CH}_2\text{CHO}$.

.....
 [1]

[Total: 21]

4 Fig. 4.1 shows some reactions of compound **D**, 2-bromobutane.




Fig. 4.1

(a) (i) State the reagent and conditions used to form **E** in reaction 1.

..... [1]

(ii) Draw the structure of **one** repeat unit of the addition polymer that forms from **E**.

[1]

(iii) **E** also forms when **F** is heated strongly in the presence of an Al_2O_3 catalyst.

Write an equation for this reaction.

..... [1]

(b) (i) Predict what is observed in reaction 2.

..... [1]

(ii) Identify the yellow precipitate and the organic ion formed in reaction 3.

yellow precipitate

organic ion [2]

(c) (i) State the type of reaction that occurs in reaction 4.

..... [1]

(ii) Reaction 5 is similar to the reaction of LiAlH_4 with carboxylic acids to form alcohols.

Suggest the role of LiAlH_4 in reaction 5.

..... [1]

(d) (i) Fig. 4.2 shows the infrared spectrum of one of the compounds **D**, **E**, **F**, **G** or **H**.




Fig. 4.2

Use information from Table 4.1 (on page 14) to identify which of the compounds **D**, **E**, **F**, **G** or **H** produces the infrared spectrum in Fig. 4.2.

Explain your answer.

.....

 [2]

Table 4.1

bond	functional groups containing the bond	characteristic infrared absorption range (in wavenumbers) / cm^{-1}
C–O	hydroxy, ester	1040–1300
C=C	aromatic compound, alkene	1500–1680
C=O	amide carbonyl, carboxyl ester	1640–1690 1670–1740 1710–1750
C≡N	nitrile	2200–2250
C–H	alkane	2850–2950
N–H	amine, amide	3300–3500
O–H	carboxyl hydroxy	2500–3000 3200–3600

- (ii) In the mass spectrum of **D**, the relative abundance of the molecular ion peak is 3.4.

Predict the relative abundance of the M+2 peak for **D**.

Explain your answer.

.....

.....

..... [1]

[Total: 11]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)

The Periodic Table of Elements

Group																																									
1	2													13	14	15	16	17	18																						
1 H hydrogen 1.0																																									
Key atomic number atomic symbol name relative atomic mass																																									
3	4													5	6	7	8	9	10	11	12	13	14	15	16	17	18														
Li	Be													B	C	N	O	F	Ne																						
lithium	beryllium													boron	carbon	nitrogen	oxygen	fluorine	neon																						
6.9	9.0													10.8	12.0	14.0	16.0	19.0	20.2																						
11	12													13	14	15	16	17	18																						
Na	Mg													Al	Si	P	S	Cl	Ar																						
sodium	magnesium													aluminium	silicon	phosphorus	sulfur	chlorine	argon																						
23.0	24.3													27.0	28.1	31.0	32.1	35.5	39.9																						
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54						
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe						
potassium	calcium	scandium	titanium	vanadium	chromium	manganese	iron	cobalt	nickel	copper	zinc	gallium	germanium	arsenic	selenium	bromine	krypton	rubidium	strontium	yttrium	zirconium	niobium	molybdenum	technetium	ruthenium	rhodium	palladium	silver	cadmium	indium	tin	antimony	tellurium	iodine	xenon						
39.1	40.1	45.0	47.9	50.9	52.0	54.9	55.8	58.9	58.7	63.5	65.4	69.7	72.6	74.9	79.0	79.9	83.8	85.5	87.6	88.9	91.2	92.9	95.9	101.1	102.9	106.4	107.9	112.4	114.8	118.7	127.6	126.9	131.3	131.3							
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86						
Rb	Sr	lanthanoids														Po	At	Rn	Ba	La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu							
rubidium	strontium															polonium	astatine	radon	barium	cerium	praseodymium	neodymium	promethium	samarium	europium	gadolinium	terbium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium								
85.5	87.6															—	—	—	137.3	140.1	140.9	144.4	—	150.4	152.0	157.3	158.9	162.5	164.9	167.3	173.1	175.0	175.0								
55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89–103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118						
Cs	Ba	actinoids														Tl	Pb	Bi	Po	At	Rn	Fr	Ra	actinoids																	
caesium	barium															thallium	lead	bismuth	polonium	astatine	radon	francium	radium																		
132.9	137.3															204.4	207.2	209.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
87	88	89–103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	87	88	89–103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118						
Fr	Ra	actinoids														Nh	Fl	Mc	Lv	Ts	Og	Fr	Ra	actinoids																	
francium	radium															nihonium	flerovium	moscovium	livermorium	tennessine	oganesson	francium	radium																		

lanthanoids

57La138.9lanthanum

62Sm150.4samarium

63Eu152.0europium

64Gd157.3gadolinium

65Tb158.9terbium

66Dy162.5dysprosium

67Ho164.9holmium

68Er167.3erbium

71Lu175.0lutetium

actinoids

89Ac231.0actinium

90Th232.0thorium

91Pa231.0protactinium

92U238.0uranium

93Np237.0neptunium

94Pu244.0plutonium

95Am243.0americium

96Cm247.0curium

97Bk247.0berkelium

98Cf251.0californium

99Es261.0einsteinium

100Fm267.0fermium

101Md288.1mendelevium

102No289.1nobelium

103Lr260.1lawrencium

lanthanoids

actinoids