



Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/31

Paper 3 Advanced Practical Skills 1

May/June 2026

2 hours

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.
- Notes for use in qualitative analysis are provided in the question paper.

Session

Laboratory

For Examiner's Use

1	
2	
3	
Total	

This document has **12** pages.

Qualitative analysis

For each test, you should record all your observations in the spaces provided.

Examples of observations include:

- colour changes seen
- the formation of any precipitate and its solubility (where appropriate) in an excess of the reagent added
- the formation of any gas and its identification (where appropriate) by a suitable test.

You should record clearly at what stage in a test an observation is made.

Where no change is observed, you should write 'no change'.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

If any solution is warmed, a boiling tube must be used. If a solid is heated, a hard-glass test-tube must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

No additional tests should be attempted.

1 (a) FA 1 is a solution containing one anion and one cation, both of which are listed in the Qualitative analysis notes.

- (i) Carry out the following tests and record your observations in Tables 1.1 and 1.2. For each test use a 1 cm depth of **FA 1** in a test-tube.

Table 1.1

test	observations
Test 1 Add aqueous sodium hydroxide, then	
add a few drops of acidified aqueous potassium manganate(VII) and shake the tube, then	
add a few drops of aqueous hydrogen peroxide.	
Test 2 Add aqueous ammonia.	



Table 1.2

test	observations
Test 3 Add a few drops of aqueous silver nitrate, then ----- add aqueous ammonia.	
Test 4 Add 1 cm depth of aqueous sodium carbonate, then ----- add dilute nitric acid.	

[6]

(ii) Deduce the formulae of the ions present in **FA 1**.
 cation anion
 [1]
(iii) Write an ionic equation for **one** of the reactions in Test 4. Include state symbols.

..... [1]

(iv) Explain why the addition of aqueous ammonia in Test 3 does **not** help you to identify the anion.

 [1]



(b) **FA 2** is a solution containing one cation and one anion, both of which are listed in the Qualitative analysis notes. The anion contains only elements in Group 16 of the Periodic Table.

- (i) Carry out tests to identify both of the ions in **FA 2**.
Record your tests and observations in a suitable form.

[5]

- (ii) Deduce the formula of **FA 2**.

FA 2 [1]

[Total: 15]



Quantitative analysis

Read through the whole method before starting any practical work. Where appropriate, prepare a table for your results in the space provided.

Show the precision of the apparatus you used in the data you record.

Show your working and appropriate significant figures in the answer to **each** step of your calculations.

2 FA 3 is a hydrated salt. **FA 3** was dissolved in water to prepare the **FA 2** used in **Question 1**.

FA 3 decomposes when heated, losing its water of crystallisation.

You will determine the formula of **FA 3** by heating it until it becomes anhydrous.

(a) Method

- Weigh the empty crucible with its lid. Record the mass in the results section.
- Add 2.10–2.40 g of **FA 3** to the crucible.
- Weigh the crucible and lid with **FA 3**. Record the mass.
- Place the crucible on the pipeclay triangle. Heat the crucible and contents **gently**, with the lid on, for approximately 2 minutes.
- Remove the lid. Heat the crucible and contents strongly for approximately 5 minutes.
- Replace the lid and leave the crucible to cool for at least 5 minutes.

While the crucible is cooling, you may wish to begin work on Question 3.

- When the crucible is cool, weigh the crucible and contents with the lid on. Record the mass.
- Place the crucible on the pipeclay triangle. Remove the lid.
- Heat the crucible and contents strongly for a further 2 minutes.
- Replace the lid and leave the crucible to cool for at least 5 minutes.
- When the crucible is cool, reweigh the crucible and contents with the lid on. Record the mass.
- Calculate the mass of **FA 3** added to the crucible, the mass of residue obtained and the mass loss. Record these masses.

Results

I	
II	
III	
IV	

[4]

(b) Calculations

- (i) Calculate the amount, in mol, of water of crystallisation lost during the decomposition of **FA 3**.

amount of H_2O lost = mol [1]

- (ii) Use the formula of **FA 2** in **1(b)(ii)** to calculate the relative formula mass, M_r , of the anhydrous salt of **FA 3**.

(If you were unable to identify **FA 2** in **1(b)(ii)**, then assume it is barium nitrate. This is **not** the correct identity.)

M_r of anhydrous salt of **FA 3** = [1]

- (iii) Use your answer to **2(b)(ii)** to calculate the amount, in mol, of the anhydrous salt of **FA 3** produced by decomposition in your experiment in **2(a)**.

amount of anhydrous salt of **FA 3** = mol [1]

- (iv) Deduce the formula of the hydrated salt, **FA 3**.
Show your working.

FA 3 [1]

- (c) (i) State the uncertainty in a single reading for the balance that you used.

uncertainty = $\pm$ g

Calculate the maximum percentage error in your measurement of the mass of **FA 3**.
Show your working.

maximum percentage error = % [1]



- (ii) A student suggests that the experiment in **2(a)** will be more accurate if the mass of **FA 3** used is doubled.

Give **one** reason why the student's suggestion is **not** correct.

.....

.....

..... [1]

[Total: 10]

- 3 You will determine the concentration of a solution of sulfuric acid by titration with aqueous potassium hydroxide.

FA 4 is dilute sulfuric acid, H_2SO_4 .

FA 5 is aqueous potassium hydroxide containing 10.0 g dm^{-3} KOH.

FA 6 is bromophenol blue indicator.

(a) Method

Preparation of **FA 7**

- Pipette 25.0 cm^3 of **FA 4** into the 250 cm^3 volumetric flask.
- Make the solution up to 250 cm^3 with distilled water.
- Thoroughly mix the contents of the volumetric flask.
- This solution of sulfuric acid is **FA 7**.

Titration

- Fill the burette with **FA 7**.
- Rinse the pipette with distilled water and then with **FA 5**.
- Pipette 25.0 cm^3 of **FA 5** into a conical flask.
- Add a few drops of **FA 6** to the solution in the conical flask.
- Perform a rough titration and record your burette readings.

rough titre = cm^3

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record in a suitable form all your burette readings and the volume of **FA 7** added in each accurate titration.

Results

I	
II	
III	
IV	
V	
VI	
VII	

[7]



- (b) From your accurate titration results, calculate a suitable mean value to use in your calculations. Show clearly how you obtain the mean value.

25.0 cm³ of **FA 5** required cm³ of **FA 7**. [1]

(c) Calculations

- (i) Give your answers to **3(c)(ii)**, **3(c)(iv)** and **3(c)(v)** to an appropriate number of significant figures. [1]

- (ii) Calculate the amount, in mol, of potassium hydroxide in the volume of **FA 5** used in each titration.

amount of KOH = mol [1]

- (iii) Give the equation for the reaction taking place in the titration. Include state symbols.

..... [1]

- (iv) Calculate the concentration, in mol dm⁻³, of sulfuric acid in **FA 7**.

concentration of H₂SO₄ = mol dm⁻³ [1]

- (v) Calculate the mass, in g, of sulfuric acid in 1.00 dm³ of **FA 4**.

mass of H₂SO₄ = g [1]

- (d) A student carries out the procedure in **3(a)**. However, by mistake the student was given alkali containing 10.0 g dm⁻³ of sodium hydroxide instead of 10.0 g dm⁻³ of potassium hydroxide.

The student was **not** aware of this mistake. State and explain what effect the wrong alkali has on the concentration of **FA 7** that the student calculates in **3(c)(iv)**.

..... [2]

[Total: 15]



Qualitative analysis notes

Reactions of cations

cation	reaction with	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	no ppt. ammonia produced on warming	–
barium, Ba ²⁺ (aq)	faint white ppt. is observed unless [Ba ²⁺ (aq)] is very low	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. unless [Ca ²⁺ (aq)] is very low	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	pale blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

Reactions of anions

anion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, Cl ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream/off-white ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives pale yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil
nitrite, NO ₂ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil; decolourises acidified aqueous KMnO ₄
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids); gives white ppt. with high [Ca ²⁺ (aq)]
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids); decolourises acidified aqueous KMnO ₄
thiosulfate, S ₂ O ₃ ²⁻ (aq)	gives off-white/pale yellow ppt. slowly with H ⁺



Tests for gases

gas	test and test result
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater
hydrogen, H_2	'pops' with a lighted splint
oxygen, O_2	relights a glowing splint

Tests for elements

element	test and test result
iodine, I_2	gives blue-black colour on addition of starch solution

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 $\text{J g}^{-1} \text{ K}^{-1}$)

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.



The Periodic Table of Elements

Group																			
1	2	Key												13	14	15	16	17	18
		1Hhydrogen1.0																	
3Li lithium 6.9	4Be beryllium 9.0	atomic number atomic symbol name relative atomic mass												5B boron 10.8	6C carbon 12.0	7N nitrogen 14.0	8O oxygen 16.0	9F fluorine 19.0	10Ne neon 20.2
11Na sodium 23.0	12Mg magnesium 24.3													13Al aluminium 27.0	14Si silicon 28.1	15P phosphorus 31.0	16S sulfur 32.1	17Cl chlorine 35.5	18Ar argon 39.9
19K potassium 39.1	20Ca calcium 40.1	21Sc scandium 45.0	22Ti titanium 47.9	23V vanadium 50.9	24Cr chromium 52.0	25Mn manganese 54.9	26Fe iron 55.8	27Co cobalt 58.9	28Ni nickel 58.7	29Cu copper 63.5	30Zn zinc 65.4	31Ga gallium 69.7	32Ge germanium 72.6	33As arsenic 74.9	34Se selenium 79.0	35Br bromine 79.9	36Kr krypton 83.8		
37Rb rubidium 85.5	38Sr strontium 87.6	39Y yttrium 88.9	40Zr zirconium 91.2	41Nb niobium 92.9	42Mo molybdenum 95.9	43Tc technetium —	44Ru ruthenium 101.1	45Rh rhodium 102.9	46Pd palladium 106.4	47Ag silver 107.9	48Cd cadmium 112.4	49In indium 114.8	50Sn tin 118.7	51Sb antimony 121.8	52Te tellurium 127.6	53I iodine 126.9	54Xe xenon 131.3		
55Cs caesium 132.9	56Ba barium 137.3	lanthanoids		73Ta tantalum 180.9	74W tungsten 183.8	75Re rhenium 186.2	76Os osmium 190.2	77Ir iridium 192.2	78Pt platinum 195.1	79Au gold 197.0	80Hg mercury 200.6	81Tl thallium 204.4	82Pb lead 207.2	83Bi bismuth 209.0	84Po polonium —	85At astatine —	86Rn radon —		
87Fr francium —	88Ra radium —	actinoids		105Db dubnium —	106Sg seaborgium —	107Bh bohrium —	108Hs hassium —	109Mt meitnerium —	110Ds darmstadtium —	111Rg roentgenium —	112Cn copernicium —	113Nh nihonium —	114Fl flerovium —	115Mc moscovium —	116Lv livermorium —	117Ts tennessine —	118Og oganesson —		

lanthanoids

actinoids

57	La	lanthanum	138.9	58	Ce	cerium	140.1	59	Pr	praseodymium	140.9	60	Nd	neodymium	144.2	61	Pm	promethium	—	62	Sm	samarium	150.4	63	Eu	euporium	152.0	64	Gd	gadolinium	157.3	65	Tb	terbium	158.9	66	Dy	dyprosium	162.5	67	Ho	holmium	164.9	68	Er	erbium	167.3	69	Tm	thulium	168.9	70	Yb	yterbium	173.1	71	Lu	lutetium	175.0
89	Ac	actinium	—	90	Th	thorium	232.0	91	Pa	protactinium	231.0	92	U	uranium	238.0	93	Np	neptunium	—	94	Pu	plutonium	—	95	Am	americium	—	96	Cm	curium	—	97	Bk	berkelium	—	98	Cf	californium	—	99	Es	einsteinium	—	100	Fm	fermium	—	101	Md	mendelevium	—	102	No	nobelium	—	103	Lr	lawrencium	—