

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/12

Paper 1 Pure Mathematics 1 (P1)

May/June 2017

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.



- 1 (i) Find the coefficient of x in the expansion of $\left(2x - \frac{1}{x}\right)^5$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Hence find the coefficient of x in the expansion of $(1 + 3x^2)\left(2x - \frac{1}{x}\right)^5$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 The point A has coordinates $(-2, 6)$. The equation of the perpendicular bisector of the line AB is $2y = 3x + 5$.

(i) Find the equation of AB .

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the coordinates of B .

[3]

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

3 (i) Prove the identity $\left(\frac{1}{\cos \theta} - \tan \theta\right)^2 \equiv \frac{1 - \sin \theta}{1 + \sin \theta}$. [3]

[illegible]

© UCLES 2017



-
- This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dashed lines spaced evenly down the page, providing a guide for letter height and placement. The background is plain white, and there are no other markings or text present.

- (ii)** In the case where $r = 5$ and $\theta = \frac{1}{6}\pi$, find the area of the shaded region. [4]

[illegible]


5 A curve has equation $y = 3 + \frac{12}{2-x}$.

(i) Find the equation of the tangent to the curve at the point where the curve crosses the x -axis. [5]

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dotted lines, providing a guide for letter height and placement. The lines are evenly spaced across the entire page, leaving ample room for writing practice. There is no text or other markings on the page.

- (ii) A point moves along the curve in such a way that the x -coordinate is increasing at a constant rate of 0.04 units per second. Find the rate of change of the y -coordinate when $x = 4$. [2]

[illegible]



The diagram shows the straight line $x + y = 5$ intersecting the curve $y = \frac{4}{x}$ at the points $A(1, 4)$ and $B(4, 1)$. Find, showing all necessary working, the volume obtained when the shaded region is rotated through 360° about the x -axis. [7]

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dashed lines, providing a guide for letter height and placement. The lines are evenly spaced across the entire page, which is otherwise blank.

- 7 (a) The first two terms of an arithmetic progression are 16 and 24. Find the least number of terms of the progression which must be taken for their sum to exceed 20 000. [4]

[illegible]

- (b) A geometric progression has a first term of 6 and a sum to infinity of 18. A new geometric progression is formed by squaring each of the terms of the original progression. Find the sum to infinity of the new progression. [4]

[illegible]

- 8** Relative to an origin O , the position vectors of three points A , B and C are given by

$$\overrightarrow{OA} = 3\mathbf{i} + p\mathbf{j} - 2p\mathbf{k}, \quad \overrightarrow{OB} = 6\mathbf{i} + (p+4)\mathbf{j} + 3\mathbf{k} \quad \text{and} \quad \overrightarrow{OC} = (p-1)\mathbf{i} + 2\mathbf{j} + q\mathbf{k},$$

where p and q are constants.

- (i) In the case where $p = 2$, use a scalar product to find angle AOB . [4]

This image shows a single page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

This image shows a full page of a handwriting practice worksheet. It consists of approximately 20 horizontal rows. Each row is defined by two parallel dotted lines, creating a series of uniform gaps for letter height. The entire page is otherwise blank, with no margins, text, or other markings.

9 The equation of a curve is $y = 8\sqrt{x} - 2x$.

(i) Find the coordinates of the stationary point of the curve. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find an expression for $\frac{d^2y}{dx^2}$ and hence, or otherwise, determine the nature of the stationary point. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find the values of x at which the line $y = 6$ meets the curve.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iv) State the set of values of k for which the line $y = k$ does not meet the curve.

[1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 The function f is defined by $f(x) = 3 \tan\left(\frac{1}{2}x\right) - 2$, for $-\frac{1}{2}\pi \leq x \leq \frac{1}{2}\pi$.

(i) Solve the equation $f(x) + 4 = 0$, giving your answer correct to 1 decimal place. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find an expression for $f^{-1}(x)$ and find the domain of f^{-1} . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Sketch, on the same diagram, the graphs of $y = f(x)$ and $y = f^{-1}(x)$.

[3]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.