Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

7756958613

MATHEMATICS **9709/41**

Paper 4 Mechanics **May/June 2022**

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use $10\,\text{m}\,\text{s}^{-2}$.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Any blank pages are indicated.

BLANK PAGE

1 A car starts from rest and moves in a straight line with constant acceleration for a distance of 200 m, reaching a speed of 25 m s^{-1}. The car then travels at this speed for 400 m, before decelerating uniformly to rest over a period of 5 s.

(a) Find the time for which the car is accelerating. [2]

.....

.....

.....

.....

.....

(b) Sketch the velocity–time graph for the motion of the car, showing the key points. [2]

(c) Find the average speed of the car during its motion. [2]

.....

.....

.....

.....

.....

.....

2 Two particles P and Q, of masses 0.5 kg and 0.3 kg respectively, are connected by a light inextensible string. The string is taut and P is vertically above Q. A force of magnitude 10 N is applied to P vertically upwards.

Find the acceleration of the particles and the tension in the string connecting them. [5]

3 A crate of mass 300 kg is at rest on rough horizontal ground. The coefficient of friction between the crate and the ground is 0.5. A force of magnitude X N, acting at an angle α above the horizontal, is applied to the crate, where $\sin \alpha = 0.28$.

Find the greatest value of X for which the crate remains at rest. [5]

4



Three coplanar forces of magnitudes 20 N, 100 N and F N act at a point. The directions of these forces are shown in the diagram.

Given that the three forces are in equilibrium, find F and α. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 Two racing cars A and B are at rest alongside each other at a point O on a straight horizontal test track. The mass of A is 1200 kg. The engine of A produces a constant driving force of 4500 N. When A arrives at a point P its speed is $25\,\text{m}\,\text{s}^{-1}$. The distance OP is d m. The work done against the resistance force experienced by A between O and P is 75 000 J.

(a) Show that $d = 100$. [3]

Car B starts off at the same instant as car A. The two cars arrive at P simultaneously and with the same speed. The engine of B produces a driving force of 3200 N and the car experiences a constant resistance to motion of 1200 N.

(b) Find the mass of B. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the steady speed which B can maintain when its engine is working at the same rate as it is at P. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 A particle starts from a point O and moves in a straight line. The velocity $v\,\text{m s}^{-1}$ of the particle at time t s after leaving O is given by

$$v = k(3t^2 - 2t^3),$$

where k is a constant.

(a) Verify that the particle returns to O when $t = 2$. [4]

(b) It is given that the acceleration of the particle is $-13.5\,\text{m s}^{-2}$ for the positive value of t at which $v = 0$.

Find k and hence find the total distance travelled in the first two seconds of motion. [6]

7 Two particles A and B, of masses 0.4 kg and 0.2 kg respectively, are moving down the same line of greatest slope of a smooth plane. The plane is inclined at 30° to the horizontal, and A is higher up the plane than B. When the particles collide, the speeds of A and B are $3\,\text{m s}^{-1}$ and $2\,\text{m s}^{-1}$ respectively. In the collision between the particles, the speed of A is reduced to $2.5\,\text{m s}^{-1}$.

(a) Find the speed of B immediately after the collision. [2]

.....

.....

.....

.....

.....

.....

.....

After the collision, when B has moved 1.6 m down the plane from the point of collision, it hits a barrier and returns back up the same line of greatest slope. B hits the barrier 0.4 s after the collision, and when it hits the barrier, its speed is reduced by 90%. The two particles collide again 0.44 s after their previous collision, and they then coalesce on impact.

(b) Show that the speed of B immediately after it hits the barrier is $0.5\,\text{m s}^{-1}$. Hence find the speed of the combined particle immediately after the second collision between A and B. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

BLANK PAGE

BLANK PAGE