

Cambridge International AS & A Level

MATHEMATICS**9709/32**

Paper 3 Pure Mathematics 3

May/June 2026**1 hour 50 minutes**

You must answer on the answer booklet.

You will need: Answer booklet
List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Write your answers on the answer booklet. Follow the instructions on the front cover of the answer booklet. If you need additional answer paper, ask the invigilator for a continuation booklet.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **4** pages.

1 Find the quotient and remainder when $4x^3 - 4x^2 + 7x + 2$ is divided by $2x - 1$. [3]

2 Solve the equation $2^{5-2x} = \frac{4^{x+1}}{3}$. Give your answer in the form $\frac{\ln a}{\ln b}$, where a and b are integers. [4]

3 The complex numbers s and t are given by

$$s = 3(\cos 2 + i \sin 2) \quad \text{and} \quad t = 2e^{-i}.$$

(a) Express st in the form $re^{i\theta}$, where $-\pi < \theta \leq \pi$ and $r > 0$. [2]

(b) In an Argand diagram with origin O , the points A and B represent the complex numbers s and st respectively.

By considering the line segments OA and OB , or otherwise, state the two geometric effects on OA of multiplying s by t . [2]

4 The square roots of $3 - (6\sqrt{2})i$ can be expressed in the Cartesian form $x + iy$, where x and y are real and exact.

By first forming a quartic equation, find the square roots of $3 - (6\sqrt{2})i$. Give your answer in exact Cartesian form. [5]

5 The parametric equations of a curve are

$$x = 2\theta - \tan 2\theta, \quad y = 6\theta + 2 \ln(\cos 2\theta),$$

for $0 \leq \theta \leq \frac{1}{4}\pi$.

(a) Find an expression for $\frac{dy}{dx}$ in terms of $\tan 2\theta$. [6]

(b) Find the exact coordinates of the point on the curve at which the gradient of the tangent is equal to -1 . [4]

6 The equation of a curve is $y = x \tan^{-1} 3x$ for $-\pi \leq x \leq \pi$.

(a) Find the exact value of the gradient of the curve when $x = \frac{1}{3}$. [4]

(b) Find the exact value of $\int_0^{\frac{1}{3}} y \, dx$. [6]

- 7 The equation of a curve is

$$y = \frac{3 + \sin 3x}{2 - \cos 3x}$$

for $0 \leq x \leq \pi$. The gradient of the curve is equal to $-\frac{3}{2}$ when $x = a$.

- (a) Show that a satisfies the equation $\sin^2 3a + 6 \sin 3a - 3 = 0$. [4]

- (b) Hence find the values of a for $0 \leq a \leq \pi$. [3]

- 8 (a) By sketching a suitable pair of graphs, show that the equation $\operatorname{cosec} \frac{1}{2}x = \frac{1}{2}(x+3)$ has exactly one root in the interval $0 \leq x \leq \pi$. [2]

- (b) Show by calculation that this root lies between 1 and 1.2. [2]

- (c) Show that, if a sequence of real values given by the iterative formula

$$x_{n+1} = 2 \sin^{-1} \left(\frac{2}{x_n + 3} \right)$$

converges, then it converges to the root of the equation in 8(a). [2]

- (d) Use the iterative formula in 8(c) to calculate this root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

- 9 The points A and B have position vectors $\begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 4 \\ 3 \\ 1 \end{pmatrix}$ respectively.

The line m passes through the points A and B .

- (a) Find a vector equation for m . [2]

The line l has equation $\mathbf{r} = \begin{pmatrix} 6 \\ -1 \\ -3 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -1 \\ -2 \end{pmatrix}$.

- (b) Determine whether the lines l and m are parallel, intersect or are skew. [4]

- (c) The point P lies on l and is such that angle $PAB = 60^\circ$.

Find the position vector of P . [6]

Question 10 is on the back page.

10 (a) Express $\frac{x+3}{(x-1)(2x-6)}$ in partial fractions. [4]

(b) The variables x and y satisfy the differential equation

$$(x-1)(2x-6)\frac{dy}{dx} = \frac{x+3}{\ln(2y)}.$$

It is given that $y = 1$ when $x = 4$.

Solve the differential equation to obtain an equation in x and y . [7]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.