

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/12

Paper 1 Pure Mathematics 1

October/November 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

BLANK PAGE

- 1** The coefficient of x^3 in the expansion of $(1 + kx)(1 - 2x)^5$ is 20.

Find the value of the constant k .

[4]

[illegible]

- 2** The first, second and third terms of a geometric progression are $2p + 6$, $-2p$ and $p + 2$ respectively, where p is positive.

Find the sum to infinity of the progression.

[5]

This image shows a single page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

- 3** The equation of a curve is $y = 2x^2 + m(2x + 1)$, where m is a constant, and the equation of a line is $y = 6x + 4$.

Show that, for all values of m , the line intersects the curve at two distinct points. [5]

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

- 4 The sum, S_n , of the first n terms of an arithmetic progression is given by

$$S_n = n^2 + 4n.$$

The k th term in the progression is greater than 200.

Find the smallest possible value of k .

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 Functions f and g are defined by

$$f(x) = 4x - 2, \quad \text{for } x \in \mathbb{R},$$

$$g(x) = \frac{4}{x+1}, \quad \text{for } x \in \mathbb{R}, x \neq -1.$$

(a) Find the value of $fg(7)$.

[1]

.....

.....

.....

.....

.....

(b) Find the values of x for which $f^{-1}(x) = g^{-1}(x)$.

[5]

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

- 6 (a) Prove the identity $\left(\frac{1}{\cos x} - \tan x\right)\left(\frac{1}{\sin x} + 1\right) \equiv \frac{1}{\tan x}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence solve the equation $\left(\frac{1}{\cos x} - \tan x\right)\left(\frac{1}{\sin x} + 1\right) = 2 \tan^2 x$ for $0^\circ \leq x \leq 180^\circ$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 The point (4, 7) lies on the curve $y = f(x)$ and it is given that $f'(x) = 6x^{-\frac{1}{2}} - 4x^{-\frac{3}{2}}$.

- (a) A point moves along the curve in such a way that the x -coordinate is increasing at a constant rate of 0.12 units per second.

Find the rate of increase of the y -coordinate when $x = 4$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find the equation of the curve.

[4]

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



- [illegible]

- (b)** In the case where $r = 10$ and $\theta = 0.6$, find the perimeter of the shaded region. [4]

[illegible]

- 9 A circle has centre at the point $B(5, 1)$. The point $A(-1, -2)$ lies on the circle.

(a) Find the equation of the circle. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Point C is such that AC is a diameter of the circle. Point D has coordinates $(5, 16)$.

(b) Show that DC is a tangent to the circle. [4]

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

[2]

[illegible]



(a) Find expressions for $\frac{dy}{dx}$, $\frac{d^2y}{dx^2}$ and $\int y dx$. [6]

[illegible]

- (b) Find, by calculation, the x -coordinate of M . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Find the area of the shaded region bounded by the curve and the coordinate axes. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

11 A curve has equation $y = 3 \cos 2x + 2$ for $0 \leq x \leq \pi$.

(a) State the greatest and least values of y . [2]

.....

.....

.....

.....

(b) Sketch the graph of $y = 3 \cos 2x + 2$ for $0 \leq x \leq \pi$. [2]

(c) By considering the straight line $y = kx$, where k is a constant, state the number of solutions of the equation $3 \cos 2x + 2 = kx$ for $0 \leq x \leq \pi$ in each of the following cases.

(i) $k = -3$ [1]

.....

.....

(ii) $k = 1$ [1]

.....

.....

(iii) $k = 3$ [1]

.....

.....

Functions f , g and h are defined for $x \in \mathbb{R}$ by

$$f(x) = 3 \cos 2x + 2,$$

$$g(x) = f(2x) + 4,$$

$$h(x) = 2f\left(x + \frac{1}{2}\pi\right).$$

- (d) Describe fully a sequence of transformations that maps the graph of $y = f(x)$ on to $y = g(x)$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (e) Describe fully a sequence of transformations that maps the graph of $y = f(x)$ on to $y = h(x)$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.