



Cambridge International AS & A Level

MATHEMATICS

9709/13

Paper 1 Pure Mathematics 1

October/November 2021

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2021 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **15** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1	{Reflection} {[in the] x -axis} or {Stretch of scale factor -1} {parallel to y -axis}	*B1 DB1	{ } indicate how the B1 marks should be awarded throughout.
	Then {Translation} $\left\{\begin{pmatrix} 0 \\ 3 \end{pmatrix}\right\}$	B1 B1	Or Translation 3 units in the positive y -direction. N.B. If order reversed a maximum of 3 out of 4 marks awarded.
	Alternative method for question 1		
	{Translation} $\left\{\begin{pmatrix} 0 \\ -3 \end{pmatrix}\right\}$	B1 B1	Or Translation 3 units in the negative y -direction.
	Then {Reflection} {in the x -axis} or {Stretch of scale factor -1} {parallel to y -axis}	*B1 DB1	N.B. If order reversed a maximum of 3 out of 4 marks awarded.
		4	

PUBLISHED

Question	Answer	Marks	Guidance
2(a)	$1 + 6ax + 15a^2x^2$	B1	Terms must be evaluated.
		1	
2(b)	<i>their</i> $15a^2 \pm (3 \times \text{their } 6a)$	*M1	Expect $15a^2 - 18a$.
	$15a^2 - 18a = -3$	A1	
	$(3)(a-1)(5a-1) [= 0]$	DM1	Dependent on 3-term quadratic. Or solve using formula or completing the square.
	$a = 1, \frac{1}{5}$	A1	WWW. If DM0 awarded SC B1 if both answers correct.
		4	


PUBLISHED

Question	Answer	Marks	Guidance
3(a)	$\{5(y-3)^2\} \quad \{+5\}$	B1 B1	Accept $a = -3, b = 5$
		2	
3(b)	$[f'(x) =] 5x^4 - 30x^2 + 50$	B1	
	$5(x^2 - 3)^2 + 5$ or $b^2 < 4ac$ and at least one value of $f'(x) > 0$	M1	
	> 0 and increasing	A1	WWW
		3	

Question	Answer	Marks	Guidance
4(a)	$84 - 3(n-1) = 0$	M1	OE, SOI. Allow either $= 0$ or < 0 (to -3).
	Smallest n is 30	A1	SC B2 for answer only $n = 30$ WWW.
		2	
4(b)	$\left(\frac{2k}{2}\right)[168 + (2k-1)(-3)] = \left(\frac{k}{2}\right)[168 + (k-1)(-3)]$	M1 A1	M1 for forming an equation using correct formula. A1 for at least one side correct.
	$k = 19$	A1	
		3	

PUBLISHED

Question	Answer	Marks	Guidance
5(a)	$\text{Angle } XYZ = \sin^{-1}\left(\frac{9}{11}\right) = 0.9582$ or $\sin XYZ = \frac{9}{11}$ leading to $XYZ = 0.9582$	B1	AG. OE using cosine rule.
		1	
5(b)	$XY = \sqrt{11^2 - 9^2} = \sqrt{40}$ or using 0.9582 and trigonometry	*M1 A1	
	$AB = 9 + 11 - \text{their } XY$	B1 FT	OE e.g. $20 - 2\sqrt{10}$, $2 + 9 - 2\sqrt{10} + 11 - 2\sqrt{10}$
	$\text{Arc } AC = 11 \times 0.9582$	M1	
	$\text{Arc } BC = 9 \times \frac{\pi}{2}$	M1	
	$\text{Perimeter} = [13.6(8) + 10.5(4) + 14.1(4) =] 38.4$	A1	AWRT. Answer must be evaluated as a single decimal.
		6	

Question	Answer	Marks	Guidance
6(a)		B1	A reflection of the given curve in $y = x$ (the line $y = x$ can be implied by position of curve).
		1	

PUBLISHED

Question	Answer	Marks	Guidance
6(b)	$y = \frac{-x}{\sqrt{4-x^2}}$ leading to $x^2 = y^2(4-x^2)$	*M1	Squaring and clearing the fraction. Condone one error in squaring $-x$ or y
	$x^2(1+y^2) = 4y^2$	DM1	OE. Factorisation of the new subject with order of operations correct. Condone sign errors.
	$x = (\pm) \frac{2y}{\sqrt{1+y^2}}$	DM1	$x = (\pm) \sqrt{\frac{4y^2}{1+y^2}}$ OE is acceptable for this mark. Isolating the new subject. Order of operations correct. Condone sign errors.
	$f^{-1}(x) = \frac{-2x}{\sqrt{1+x^2}}$	A1	Selecting the correct square root. Must not have fractions in numerator or denominator.
		4	
6(c)	1 or $a=1$	B1	Do not allow $x=1$ or $-1 < x < 1$
		1	
6(d)	$[fg(x) = f(2x)] \frac{-2x}{\sqrt{4-4x^2}}$	B1	Allow $\frac{-2x}{\sqrt{4-(2x)^2}}$ or any correct unsimplified form.
	$fg(x) = \frac{-x}{\sqrt{1-x^2}}$ or $\frac{-x}{1-x^2} \sqrt{1-x^2}$ or $\frac{x}{x^2-1} \sqrt{1-x^2}$	B1	Result of cancelling 2 in numerator and denominator.
		2	

PUBLISHED

Question	Answer	Marks	Guidance
7(a)	$\tan x + \cos x = k(\tan x - \cos x)$ leading to $\sin x + \cos^2 x = k(\sin x - \cos^2 x)$	M1	Use $\tan x = \frac{\sin x}{\cos x}$ and clear fraction.
	$\sin x + 1 - \sin^2 x = k \sin x - k + k \sin^2 x$	*M1	Use $\cos^2 x = 1 - \sin^2 x$ twice to obtain an equation in sine.
	$k \sin^2 x + \sin^2 x + k \sin x - \sin x - k - 1 = 0$	DM1	Gather like terms on one side of the equation.
	$(k+1)\sin^2 x + (k-1)\sin x - (k+1) = 0$	A1	AG. Factorise to obtain answer.
		4	
7(b)	$5\sin^2 x + 3\sin x - 5 = 0$	B1	
	$\sin x = \frac{-3 \pm \sqrt{9+100}}{10}$	M1	Use formula or complete the square.
	$x = 48.1^\circ, 131.9^\circ$	A1 A1 FT	AWRT. Maximum A1 if extra solutions in range. FT for $180 - \text{their answer}$ or $540 - \text{their answer}$ if $\sin x$ is negative If M0 given and correct answers only SCB1B1 available. If answers in radians; 0.839, 2.30 can score SCB1 for both.
		4	

PUBLISHED

Question	Answer	Marks	Guidance
8(a)	$\int \left(\frac{5}{2} - x^{\frac{1}{2}} - x^{-\frac{1}{2}} \right) dx$	M1	OR as 2 separate integrals $\int \left(\frac{5}{2} - x^{1/2} \right) dx - \int (x^{-1/2}) dx$
	$\left\{ \frac{5}{2}x - \frac{2}{3}x^{\frac{3}{2}} \right\} \{-\} \left\{ 2x^{\frac{1}{2}} \right\}$	A1 A1 A1	If two separate integrals with no subtraction SC B1 for each correct integral.
	$\left(10 - \frac{16}{3} - 4 \right) - \left(\frac{5}{8} - \frac{1}{12} - 1 \right)$	DM1	Substitute limits $\frac{1}{4} \rightarrow 4$ at least once, must be seen.
	$\frac{9}{8}$ or 1.125	A1	WWW. Cannot be awarded if π appears in any integral.
		6	
8(b)	$\left[\frac{dy}{dx} = \right] -\frac{1}{2}x^{-\frac{3}{2}}$	B1	
	When $x = 1$, $m = -\frac{1}{2}$	M1	Substitute $x = 1$ into a differential.
	[Equation of normal is] $y - 1 = 2(x - 1)$	M1	Through (1, 1) with gradient $-\frac{1}{m}$ or $\frac{1-p}{1} = 2$
	[When $x = 0$,] $p = -1$	A1	WWW
		4	

PUBLISHED

Question	Answer	Marks	Guidance
9(a)	$x^2 + (2x + 5)^2 = 20$ leading to $x^2 + 4x^2 + 20x + 25 = 20$	M1	Substitute $y = 2x + 5$ and expand bracket.
	$(5)(x^2 + 4x + 1)[= 0]$	A1	3-term quadratic.
	$x = \frac{-4 \pm \sqrt{16 - 4}}{2}$	M1	OE. Apply formula or complete the square.
	$A = (-2 + \sqrt{3}, 1 + 2\sqrt{3})$	A1	Or 2 correct x values.
	$B = (-2 - \sqrt{3}, 1 - 2\sqrt{3})$	A1	Or all values correct. SC B1 all 4 values correct in surd form without working. SC B1 all 4 values correct in decimal form from correct formula or completion of the square
	$AB^2 = \text{their}(x_2 - x_1)^2 + \text{their}(y_2 - y_1)^2$	M1	Using <i>their</i> coordinates in a correct distance formula. Condone one sign error in $x_2 - x_1$ or $y_2 - y_1$
	$[AB^2 = 48 + 12 \text{ leading to}] AB = \sqrt{60}$	A1	OE. CAO. Do not accept decimal answer. Answer must come from use of surd form in distance formula.
		7	

PUBLISHED

Question	Answer	Marks	Guidance
9(b)	$x^2 + m^2(x - 10)^2 = 20$	*M1	Finding equation of tangent and substituting into circle equation.
	$x^2(m^2 + 1) - 20m^2x + 20(5m^2 - 1) [= 0]$	DM1	OE. Brackets expanded and all terms collected on one side of the equation.
	$[b^2 - 4ac =] 400m^4 - 80(m^2 + 1)(5m^2 - 1)$	M1	Using correct coefficients from <i>their</i> quadratic equation.
	$400m^4 - 80(5m^4 + 4m^2 - 1) = 0 \rightarrow (-80)(4m^2 - 1) = 0$	A1	OE. Must have ‘=0’ for A1.
	$m = \pm \frac{1}{2}$	A1	
	Alternative method for question 9(b)		
	Length, l of tangent, is given by $l^2 = 10^2 - 20$	M1	
	$l = \sqrt{80}$	A1	
	$\tan \alpha = \frac{\sqrt{20}}{\sqrt{80}} = \frac{1}{2}$	M1 A1	Where α is the angle between the tangent and the x -axis.
	$m = \pm \frac{1}{2}$	A1	
		5	

PUBLISHED

Question	Answer	Marks	Guidance
10(a)	$f''(x) = -\left(\frac{1}{2}x + k\right)^{-3}$	B1	
	$f''(2) > 0 \Rightarrow -(1+k)^{-3} > 0$	M1	Allow for solving <i>their</i> $f''(2) > 0$
	$k < -1$	A1	WWW
		3	
10(b)	$\left[f(x) = \int \left(\left(\frac{1}{2}x - 3 \right)^{-2} - (-2)^{-2} \right) dx = \right] \left\{ \frac{\left(\frac{1}{2}x - 3 \right)^{-1}}{-1 \times \frac{1}{2}} \right\} \left\{ -\frac{x}{4} \right\}$	B1 B1	Allow $-2 \left(\frac{1}{2}x + k \right)^{-1}$ OE for 1 st B1 and $-(1+k)^{-2}x$ OE for 2 nd B1
	$3\frac{1}{2} = 1 - \frac{1}{2} + c$	M1	Substitute $x = 2, y = 3\frac{1}{2}$ into <i>their</i> integral with c present.
	$f(x) = \frac{-2}{\left(\frac{1}{2}x - 3 \right)} - \frac{x}{4} + 3$	A1	OE
		4	
10(c)	$\left(\frac{1}{2}x - 3 \right)^{-2} - (-2)^{-2} = 0$	M1	Substitute $k = -3$ and set to zero.
	leading to $\left(\frac{1}{2}x - 3 \right)^2 = 4 \quad \left[\frac{1}{2}x - 3 = (\pm)2 \right]$ leading to $x = 10$	A1	
	$(10, -\frac{1}{2})$	A1	Or when $x = 10, y = -1 - 2\frac{1}{2} + 3 = -\frac{1}{2}$
	$f''(10) \left[= -(5-3)^{-3} \rightarrow \right] < 0 \rightarrow \text{MAXIMUM}$	A1	WWW
		4	