

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/41

Paper 4 Mechanics

October/November 2021

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 m s^{-2} .

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages.

- 1** A bus moves from rest with constant acceleration for 12 s. It then moves with constant speed for 30 s before decelerating uniformly to rest in a further 6 s. The total distance travelled is 585 m.

(a) Find the constant speed of the bus.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the magnitude of the deceleration.

[1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 Two small smooth spheres A and B , of equal radii and of masses km kg and m kg respectively, where $k > 1$, are free to move on a smooth horizontal plane. A is moving towards B with speed 6 m s^{-1} and B is moving towards A with speed 2 m s^{-1} . After the collision A and B coalesce and move with speed 4 m s^{-1} .

(a) Find k . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find, in terms of m , the loss of kinetic energy due to the collision. [2]

.....

.....

.....

.....


.....

.....

.....

.....

.....



Given that $\sin \alpha = \frac{3}{5}$, find the values of P and θ . [6]

[illegible]

- 4 A particle of mass 12 kg is stationary on a rough plane inclined at an angle of 25° to the horizontal. A force of magnitude P N acting parallel to a line of greatest slope of the plane is used to prevent the particle sliding down the plane. The coefficient of friction between the particle and the plane is 0.35.

(a) Draw a sketch showing the forces acting on the particle. [1]

(b) Find the least possible value of P . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 5** A car of mass 1600 kg travels at constant speed 20 m s^{-1} up a straight road inclined at an angle of $\sin^{-1} 0.12$ to the horizontal.

(a) Find the change in potential energy of the car in 30 s. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Given that the total work done by the engine of the car in this time is 1960 kJ, find the constant force resisting the motion. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Calculate, in kW, the power developed by the engine of the car.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Given that this power is suddenly decreased by 15%, find the instantaneous deceleration of the car.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 6** A particle P moves in a straight line starting from a point O and comes to rest 14 s later. At time t s after leaving O , the velocity $v \text{ m s}^{-1}$ of P is given by

$$v = pt^2 - qt \quad 0 \leq t \leq 6,$$

$$v = 63 - 4.5t \quad 6 \leq t \leq 14,$$

where p and q are positive constants.

The acceleration of P is zero when $t = 2$.

- (a)** Given that there are no instantaneous changes in velocity, find p and q . [3]

.....

.....

.....

.....

.....


.....

.....

.....

- (b)** Sketch the velocity-time graph. [3]

[Turn over



Two particles A and B of masses 2 kg and 3 kg respectively are connected by a light inextensible string. Particle B is on a smooth fixed plane which is at an angle of 18° to horizontal ground. The string passes over a fixed smooth pulley at the top of the plane. Particle A hangs vertically below the pulley and is 0.45 m above the ground (see diagram). The system is released from rest with the string taut. When A reaches the ground, the string breaks.

Find the total distance travelled by B before coming to instantaneous rest. You may assume that B does not reach the pulley. [8]

This image shows a full page of white paper designed for handwriting practice. It features 18 evenly spaced, horizontal dashed lines that run across the entire width of the page. The lines are thin and black, providing a guide for letter height and placement. There are no margins, text, or other markings on the page.

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.