

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

October/November 2023

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

BLANK PAGE

- 1 (a) Sketch the graph of $y = |4x - 2|$.

[1]

- (b) Solve the inequality $1 + 3x < |4x - 2|$.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2 The parametric equations of a curve are

$$x = (\ln t)^2, \quad y = e^{2-t^2},$$

for $t > 0$.

Find the gradient of the curve at the point where $t = e$, simplifying your answer.

[4]

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

- 3 The polynomial $2x^3 + ax^2 - 11x + b$ is denoted by $p(x)$. It is given that $p(x)$ is divisible by $(2x - 1)$ and that when $p(x)$ is divided by $(x + 1)$ the remainder is 12.

Find the values of a and b .

[5]

[illegible]

- 4 (a) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 4 - 3i| \leq 2$ and $\operatorname{Re} z \leq 3$. [4]

- (b) Find the greatest value of $\arg z$ for points in this region. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 Find the exact value of $\int_0^6 \frac{x(x+1)}{x^2+4} dx$.

[6]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- 6 (a)** By sketching a suitable pair of graphs, show that the equation

$$\cot x = 2 - \cos x$$

has one root in the interval $0 < x \leq \frac{1}{2}\pi$.

[2]

- (b)** Show by calculation that this root lies between 0.6 and 0.8.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[illegible]

- 7 (a)** By expressing 3θ as $2\theta + \theta$, prove the identity $\cos 3\theta \equiv 4\cos^3\theta - 3\cos\theta$. [3]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

(b) Hence solve the equation

$$\cos 3\theta + \cos \theta \cos 2\theta = \cos^2 \theta$$

for $0^\circ \leq \theta \leq 180^\circ$.

[5]

[illegible]


- 8** It is given that $\frac{2+3ai}{a+2i} = \lambda(2-i)$, where a and λ are real constants.

(a) Show that $3a^2 + 4a - 4 = 0$.

[4]

[illegible]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



-
- This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

[Turn over

10 The equations of the lines l and m are given by

$$l: \mathbf{r} = \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} \quad \text{and} \quad m: \mathbf{r} = \begin{pmatrix} 6 \\ -3 \\ 6 \end{pmatrix} + \mu \begin{pmatrix} -2 \\ 4 \\ c \end{pmatrix},$$

where c is a positive constant. It is given that the angle between l and m is 60° .

(a) Find the value of c .

[4]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- (b)** Show that the length of the perpendicular from $(6, -3, 6)$ to l is $\sqrt{11}$. [5]

[illegible]

- 11 The variables x and y satisfy the differential equation

$$x^2 \frac{dy}{dx} + y^2 + y = 0.$$

It is given that $x = 1$ when $y = 1$.

- (a) Solve the differential equation to obtain an expression for y in terms of x . [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) State what happens to the value of y when x tends to infinity. Give your answer in an exact form. [1]

.....

.....

.....

.....

.....

[illegible]

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.