

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/33

Paper 3 Pure Mathematics 3

October/November 2023

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

BLANK PAGE

- 1** Find the set of values of x satisfying the inequality $|2^{x+1} - 2| < 0.5$, giving your answer to 3 significant figures. [4]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school handwriting practice paper. The lines are evenly spaced and run across the entire width of the page. There are no margins, text, or other markings present.

- 2 On an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 1 + 2i| \leq |z|$ and $|z - 2| \leq 1$. [5]

- 3** The polynomial $2x^3 + ax^2 + bx + 6$, where a and b are constants, is denoted by $p(x)$. When $p(x)$ is divided by $(x + 2)$ the remainder is -38 and when $p(x)$ is divided by $(2x - 1)$ the remainder is $\frac{19}{2}$.

Find the values of a and b .

[5]

[illegible]

- 4 Solve the quadratic equation $(3 + i)w^2 - 2w + 3 - i = 0$, giving your answers in the form $x + iy$, where x and y are real. [5]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- 5** Find the exact coordinates of the stationary points of the curve $y = \frac{e^{3x^2-1}}{1-x^2}$. [6]

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

- 6 (a)** Show that the equation $\cot^2 \theta + 2 \cos 2\theta = 4$ can be written in the form

$$4 \sin^4 \theta + 3 \sin^2 \theta - 1 = 0. \quad [3]$$

[illegible]

[Turn over

7 The equation of a curve is $x^3 + y^2 + 3x^2 + 3y = 4$.

(a) Show that $\frac{dy}{dx} = -\frac{3x^2 + 6x}{2y + 3}$. [3]

[illegible]

- (b)** Hence find the coordinates of the points on the curve at which the tangent is parallel to the x -axis. [5]

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dotted lines, providing a guide for letter height and placement. The lines are evenly spaced across the entire page, leaving ample room for writing practice. There is no text or other markings on the page.

- 8 The variables x and y satisfy the differential equation

$$e^{4x} \frac{dy}{dx} = \cos^2 3y.$$

It is given that $y = 0$ when $x = 2$.

Solve the differential equation, obtaining an expression for y in terms of x .

[7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

9 Let $f(x) = \frac{17x^2 - 7x + 16}{(2 + 3x^2)(2 - x)}$.

(a) Express $f(x)$ in partial fractions.

[5]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- (b) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^3 . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) State the set of values of x for which the expansion in (b) is valid. Give your answer in an exact form. [1]


.....

.....

.....

.....

.....



(a) Find the equation of the tangent to the curve at the point where $x = \frac{1}{2}\pi$. [4]

[illegible]

- (b) Find the exact area of the shaded region shown in the diagram, bounded by the curve and the x -axis. [5]

[illegible]

- 11** The line l has equation $\mathbf{r} = \mathbf{i} - 2\mathbf{j} - 3\mathbf{k} + \lambda(-\mathbf{i} + \mathbf{j} + 2\mathbf{k})$. The points A and B have position vectors $-2\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ and $3\mathbf{i} - \mathbf{j} + \mathbf{k}$ respectively.

(a) Find a unit vector in the direction of l . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The line m passes through the points A and B .

(b) Find a vector equation for m . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Determine whether lines l and m are parallel, intersect or are skew. [5]

[illegible]

[illegible]

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.